# A QUARTERLY TABLE

OF THE

# Mortality

IN A

# HUNDRED AND FIFTEEN OF THE DISTRICTS

OF

# ENGLAND,

(INCLUDING THE PRINCIPAL TOWNS,)

SHEWING

THE NUMBER OF DEATHS REGISTERED IN THE SIX YEARS 1838—43, THE AVERAGE NUMBER OF DEATHS IN THE FIVE WINTERS 1838—1842;

THE DEATHS IN EACH QUARTER OF 1843;

AND THE NUMBER OF DEATHS IN THE WINTER QUARTER 1844, ENDING MARCH 31st.

PUBLISHED BY AUTHORITY OF THE REGISTRAR GENERAL



LONDON:

HARTNELL, PRINTER, RED LION COURT.

# A QUARTERLY TABLE

SHT'VO

Cillatter?

BUNDRED AND FIFTEEN OF THE DISTRICTS

# ENGLAND,

to be and the ask high restrict out put the man that he shall be all the

SEASON SON SELECTION OF THE SEASON SERVICE SER

The state of the s

MARINER HASTEROTH MET TO THE OWNER OF ESPECIALIST

: 200 705

TAUGO KOLA SAR MATZING JIRTON

### TABLE

OF THE

### MORTALITY

In 115 of the Districts of England (including the principal Towns): shewing the Number of Deaths Registered in the 6 years, the average Number of Deaths in the 5 Winters, 1838—42, and the Number of Deaths in the Winter Quarter of 1844, ending March 31st.

	HAD CA			110 1			III	-			1011, 010	0			-		-	-		1111	4
1 1 1000		1021	15.		Sinter		1	1838	-42	Vinter ng 844	-1 0.100	Na il	100	n					1838	3-42	inter
Diamarana	Popu-	De	eaths ]	Regist	ered in	the y	ears	Q nai	terly	the W endin	DISMINATION	Popu	1	Death	s Regis	tered in	the yea	ars		rterly	the W
DISTRICTS	lation	-	100	-	1484		1			2 2 2	DISTRICTS	lation	1	1	1	-	1	1	AV		E E
1000	1841	1838	1839	1840	1841	1842	1843	Of Five Years	Of Five Winters	Quart Quart March		1841	1838	1839	1840	1841	1842	1843	Of Five	Of Five	Deaths is
			1		1	1	1228	15.73	3 = 1	2010	N. Midland				1		1	los	193	THE	4
Metropolis West Districts	301326	8114	6538	6936	6599	6826	7212	1751	1952	1975	division Leicester	5093	2 117	0 129	1 173	0 135	8 1457	1255	350	394	41
North Districts	366303	1	-		30 000	8550			2370		Lincoln	3611	11 11 11	4 6 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1	A Common	186	130263	1320	17 9
Central Districts	374759	10844	9751	9108	9333	8873	9547	2395	2704	2547	Nottingham	5308	13 192 11 1	A Street	127.523	4 123	9 1429	1393	361	375	38
East Districts	393247	12434	2000	101 23	100000	9947	1000		2895	2975	Basford	5963	4 104	120	0 154	9 116	3 124	100	ALCOHOLD .	337	100
South Districts	479469	13210	11562	1	Malin		12575	3044	3346	3494	Derby	3501	5 769	83	2 92	6 91	4 909	922	217	246	26
Total	1915104	53546	46100	47030	46069	46114	49332	11943	13266	13471	N. Western division	15	11111	Dast	1 3 0%	10,1	10.00	SOT.			
S. Eastern division								-	19-11		Stockport	8567	2 2410	259	257	1 204	8 2073	2203	585	656	48
Maidstone	33210	737	726	700	648	815	774	181	203	218	Macclesfield	5601	\$ 17	1 .		141	9 1403	1551	370	431	39
Brighton	46742		898	980	1102	1126		251	255	329	Gt.Broughton,	4908	5 1040	123	7 1384	1 124	1189	1091	305	348	34
Isle of Wight	42547	655	630	737	891	721	703	182	189	207	Liverpool	223054		100	11 12	7556	7407	7458	1875	1963	199
Portsea Island	53036		1180			1152	100000	297	320	344	West Derby,	88655		61	200	35	A word		12000	1003	2 6
Windsor	23044 20502	421 381	479 367	526 402	416 395	469 394	471	116	121	170	(adj. Liverpool)	1-138		101-1	050.	18	10012	100 100	Wal.	198170	51
	20302	100	901	402	11/3	0.01	. 014		100	120	Blackburn	75091		1.	1 1 3 3 2	92	A STATE	4000	465 513	100000	50
S. Midland division	1 3		21		1-33	-			1111	Mary Par	Rochdale	60577		1	7000	7-	1	1	396	1	48
St. Albans	17051	378	310	331	300	340	291	83	85	109	Bury	77496	1 3	1990	1 0200	2064	1987	1832	506	615	61
Wycombe	34150	788	692	824	751	679	748	187	205	238	Bolton	97519	2432	2857		1	-	2596	668	772	678
Oxford	19701	359	407 687	406	450 693	414 622	359	102	112	108	Wigan	66032			A CONTRACTOR	Now 1	1		434	520	64
Northampton Bedford	28103 31767	689	678	743	714	604	762	171	199	237	Prescott	93736	1 1 5	1056	12775	1 63	1	2529	260 591	306 608	259 629
Cambridge	24453	563	547	639	659	465	597	144	156	230	Manchester	192408	4 11 44	15 0 3	1000		1-1-16	6263	1597	1764	1541
Eastern					e m				100		Salford	70228	1 1.19	2449	TO SELECT	1971	1977	2098	530	567	502
division					200		200				Ashton	173964	4488	5489	4873	4282	4786	4391	1196	1288	1244
Colchester	17790	533	437 630	488	381	417 594	393 551	113	119	136	York		103								
Norwich	25254 61846	694 1373	1852	593 1752	1362	1582	1295	150 396	164	426	division Sheffield	85076	2429	2703	2410	2231	2182	2169	598	653	290
Yarmouth	24031	417	440	504	493	516	510	118	123	196	Huddersfield	107140	1	2080		1		2451	514	594	579 607
S. Western			100	68	1103				- 13	3967	Halifax	109175	11 2 30	2402	1005.3	10 3		2459	566	645	670
division	100	1								Barrier H	Bradford	132164	3093	3251	3176	3041	3323	3344	794	858	949
Devizes	22130	426	510	481	429 456	449	638 524	115	135	148	Leeds	168667		4388	4488	4360	4615	4329	1107	1155	1088
Dorchester	23380 31333	700	362	412 808	920	437 764	836	107	219	168 262	York	41130	4	1023	1442	1206	1042	1128	289	294	327
Exeter	47105	824	646 819	888	859	886	808	214	242	247		41119	010	1217	1203	1009	1133	1040	28-2	296	272
Plymouth	36527	862	670	765	730	1141	889	208	220	269	Northern division							mole	AAF		
Redruth	48062	914	909	1011	906	1232	845	249	260	270	Sunderland	56226	1479	1524	1513	1512	1357	1247	369	387	306
Penzance	50100	762	848	976	1016	1128	996	236	246	238	Gateshead	38747	1.1	968	1016	989	909	984	238	248	216
Bath	69232	1567	1602	1974	1728	1598	1024	423	477	507	Tynemouth	55625	1112	1285	1243	1340	1327	1229	315	334	332
Western					277		200	773		2017	Newcastle-on-	71850	1829	2117	1957	2104	1726	1922	487	501	436
Bristol	64298	1842	1675	2330	1895	1771	1742	476	549	464	Carlisle	36084	855	739	901	925	840	762	213	252	271
Clifton	66233	1328	1150	1739	1519	1510	1256	362	404	425	Cockermouth	35676	10763	718	696	674	597	690	173	215	184
Stroud	38920	952	693	760	728	706	808	192	239	241	Kendal	34694	729	825	821	601	714	666	184	201	222
Chelteuham	40221	869	726	855	890	1078	824	221	250	228	Welsh division					100	0	TORE	577		33
Hereford	33646	774	687	758	688 520	862	534	188	215	244	Abergavenny	50834	1150	1206	1374	1458	1109	1116	315	373	476
Worcester	21529	568 805	514	559 638	614	586 637	641	137	189	214	Pontypool	25037		615	563	648	. 545	514	147	174	142
Kidderminster	29408	591	523	670	582	621	597	149	194		Merthyr Tydfil	52864	100	1246	1566	1423	1110	1183	343	422	569
Dudley	86028	1767	1891	2022	2296	2332	1729	515	588	549	Newtown	25958		468 823	619	539	420	477	129	143	145
Walsall	34274	619	749	913	832	927	706	202	236	197	Wrexham Holywell	39542 40787	840	791	966 864	900	738	917	217	257	266 253
Wolverhampton	80722	1556	1909	2217	2153	2319	1783	508	593	542	Anglesey	38105	645	589	632	597	624	575	154	177	201
Wolstanton' Birmingham	32669 138187	710 3359	795 3639	831 3767	757 3673	964 3579	3340	901	980	248	Total, exclusive	1161	10		201,0			1 1-11	1022	1000	
Aston	50928	1099	1058	1218	1113	1093	1081	279	288		of the metropolis	1663808	109321	116505	124664	114664	115834	13869 2	9049 3	2085	32570
Coventry	31028	684	724	616	905	890	838	191	196	234	GRAND TOTAL	6578912	162867	162605	171694	160733	161948	63201 4	0992 4	5351	46041
A COLUMN TO THE REAL PROPERTY AND ADDRESS OF THE PARTY AND ADDRESS OF T	1	-	400			-11	10 11 1	3306		1914	PERSONAL PROPERTY.	1000			NAME OF					100	
			-												12/11/01	3	OLF BUILD	CONTRACTOR OF	STREET, SQUARE,	THE REAL PROPERTY.	

<sup>\*</sup> The Deaths in the Metropolis for the Years 1840·1-2, have been derived from the Weekly Tables, the Returns for each Year comprising 52 Weeks, or 364 Days. The last quarter in the Metropolis ended March 30, 1844. The Returns from other Places are for the years ending December 31, and the quarter ending March 31st.

<sup>\*</sup> Wandsworth District is included in the return for the Metropolis.

### TABLE

Of the number of Deaths Registered in a hundred and fifteen Districts,

(including the principal Towns) in each of the four

quarters of the year 1843.

1		44	are in	PAR ME	Burning	0 6440.	A raises of agrees or a	OUT THE R	Dide.UL	1 10 1	OC DEED	Q.
100	DISTRICTS.	Population				ng	DISTRICTS.	Population			egistere	
March D	Tare of the contract of the co	1841	March 31st	June 30th	Sept.	Dec. 31st		1841	March 31st	June 30th	Sept.	Dec. 31st
	* Metropolis— West Districts North Districts Central Districts East Districts South Districts	301,326 366,303 374,759 393,247 479,469	1799 2407 2382 2718 3198	1805 2206 2304 2496 2810	1664 2017 2123 2547 2920	1944 2408 2738 3199 3647	North Midland Division—  Leicester Lincoln Nottingham	50,932 36,110 53,080	367 201 375 378	276 204 299 325	269 171 362 233	343 210 357 280
Columbia Service	Total	1,915,104	12504	11621	11271	13936	Basford	59,634 35,01 <u>5</u>	223	210		260
-	South Eastern Division—	Ha Gera	us, or	2 12700s		Jung Pa all all all all	NorthWestern Division—	- 100 OF			100	
-	Maidstone Brighton Isle of Wight Portsea Island Winchester Windsor	32,310 46,742 42,547 53,036 23,044 20,502	220 315 182 344 133 89	175 292 175 334 100 88	171 228 149 326 106 64	208 282 197 338 132 73	Stockport Macclesfield GreatBoughton(Chester) Liverpool West Derby Blackburn Preston Rochdale	223,054 88,652 75,091 77,189 60,577	700 537 310 1864 567 626 509 420	511 386 292 1913 558 562 507 405	458 320 231 1826 562 381 463 345	534 308 258 1855 625 462 459 361
	South Midland Division—  St. Albans Wycombe Oxford Northampton Bedford Cambridge		91 212 96 178 218 140	60 198 85 177 177 177 124	66 146 93 148 146 122	74 192 85 174 -221 211	Bury Bolton Wigan Prescott Chorlton Manchester Salford Ashton	77,496 97,519 66,032 43,739 93,736 192,408 70,228 173,964	530 755 416 279 551 1648 588 1314	492 676 477 248 610 1449 499 1074	371 542 416 170 710 1729 493 942	439 623 523 223 658 1437 518 1061
CONCRETE CON	Eastern Division—  Colchester Ipswich Norwich Yarmouth	17,790 25,254 61,846 24,031	104 145 378 148	99 118 339 102	76 141 300 106	114 147 278 154	York Division Sheffield Huddersfield Halifax Bradford Leeds Hull York	85,076 107,140 109,175 132,164 168,667 41,130 47,779	584 664 685 796 1136 254 266	532 779 679 938 1120 282 298	507 530 532 801 1003 276 227	546 478 563 809 1070 316 257
	Devizes Dorchester Exeter St. Thomas Plymouth Redruth Penzance Bath	22,130 23,380 31,333 47,105	186 137 214 222 274 254 283 417	194 105 143 185 206 203 250 376	121 134 184 177 182 169 229 351	132 148 295 224 227 219 234 489	Northern Division—  Sunderland Gateshead Tynemouth Newcastle-on-Tyne Carlisle Cockermouth Kendal	56,226 38,747 55,625 71,850 36,084 35,676 34,694	316 246 340 565 193 191 172	333 279 309 540 195 190 209	292 198 300 394 144 145 136	306 261 280 423 230 164 149
	Western Division  Bristol Clifton Stroud Cheltenham Hereford Shrewsbury Worcester Kidderminster Dudley Walsall Wolverhampton Wolstanton Birmingham Aston Coventry	38,920 40,221 33,646 21,529 27,130 29,408 86,028 34,274 80,722 32,669 138,187 50,928	503 381 273 241 252 148 155 150 474 215 467 271 896 263 263 265	390 294 188 197 197 117 172 155 442 192 415 220 735 277 198	331 287 155 189 173 134 144 147 399 149 422 172 713 234 183	518 394 192 197 194 135 170 145 414 150 479 179 996 307 192	Welsh Division—  Abergavenny Pontypool Merthyr Tydfil Newtown Wrexham Holywell Anglesey  TOTAL—exclusive of the Metropolis  GRAND TOTAL	50,834 25,037 52,864 25,958 39,542 40,787 38,105 4,663,808	319 176 318 130 278 259 174 31154	313 160 265 139 266 204 129 28626		271 89 372 98 208 196 144 28525

<sup>&</sup>quot; Wandsworth District is included in the return for the Metropolis.

### A TABLE OF THE MORTALITY

IN THE

### Metropolis,

### SHEWING THE NUMBER OF DEATHS FROM ALL CAUSES,

Registered in the 13 Weeks ending March 30th, 1844.

CAUSES OF DEATH	Quarterly Average of Deaths in the 5 Winters of 1839-43	Deaths in the Winter Quarter ending March 30th 1844	-	CAUSES OF DEATH	Quarterly Average of Deaths in the 5 Winters of 1839-43	Deaths in the Winter Quarter ending March 30th
ALL CAUSES	12600 +	23471	IV	-Laryngitis	9 17 19	9
Specified Causes	12550	13403		Quinsey	18	17
I.—Zymotic (or Epidemic, Ende- a mic, and Contagious) Diseases	2119	2457		Bronchitis	232	24
Sporadic Diseases:—				Pneumonia	1125	1327
II.—Dropsy, Cancer, and other Dis-				Hydrothorax	86	102
eases of uncertain or variable	1491	1282	100	Asthma	555 1822	555 1904
Seat	2034	2177		Disease of Lungs, &c	233	262
IV.—Diseases of the Lungs and of the	4101	4644	V	-Pericarditis	8	24
other Organs of Respiration ( V.—Diseases of the Heart and Blood )		40		Disease of Heart, &c	266	383
Vessels	286	416	VI	-Teething	214	- 157
VI.—Diseases of the Stomach, Liver, and other Organs of Digestion	768	795	- 10	Gastritis	} 190	19 141
VII.—Diseases of the Kidneys, &c	67	2 93	2 2	Peritonitis	2 14 3	30
VIII.—Childbirth, Diseases of the Uterus, &c.	141	114	1	Tabes Mesenterica	59	100
IX.—Rheumatism, Diseases of the	. 81	74	1	Ascites	7	21
Bones, Joints, &c	. 61	14		Ulceration (of Intestines, &c.)	18	21
X.—Diseases of the Skin, Cellular Tissue, &c	18	23	4	Hernia	27 24	34
XI.—Old Age	1110	1018	1 11.3	Intussusception	6	6
XII.—Violence, Privation, and In-	333	310	46 31	Stricture	. 8 .	6
I.—Small Pox	263	252		Hæmatemesis	49	11
Measles	239	334	1	Disease of Pancreas	.2	66
Scarlatina	309 524	536 487		Hepatitis	14	16
Croup Thrush.	105 45	107 45		Jaundice	26	28
Diarrhœa	64	79	. 0	Disease of Liver, &c	102	97
Dysentery	20 2	29	VII.	Nephritis	6 5	3
Influenza	49	66	-	Ischuria	2	1
Remittent Fever	6	6	35	Diabetes	6	. 8
Typhus Erysipelas	406	432	1 2	Cystitis	5	6 6
Syphilis	8 2	12		Stricture	5	14
II.—Inflammation	94	18		Disease of Kidneys, &c	39	55?
Hæmorrhage	38	30	VIII.	-Childbirth	104	80
Dropsy	483	392		Paramenia Ovarium Dropsy	5	7
Mortification	65	49	A. J.	Disease of Uterus, &c	29	27
Scrofula	26	36	IX	Arthritis	= .6	1
CancerTumor	94 22	141 13	1	Rheumatism	36	31
Gout	19 78	14	The Contract of the Contract o	Disease of Joints, &c	40	42
Debility	279	214	X	Carbuncle	6 .4	3
Malformations	228	23	11 25	Ulcer	8	11
III Cephalitis	140	160		Fistula	5	6
Hydrocephalus	256	481 301	2 2	Disease of Skin, &c	2 4	3
Paralysis	228	281,	15 16	-Old Age	1110	1018
Convulsions Tetanus	739	8	XII	-Intemperance	6	7
Chorea Epilepsy	62	3 57	75 2	Privation	317	292
Insanity	15	22	22 .52	Causes not specified	e 50	68
Delirium Tremens Disease of Brain, &c	18 124	143		causes not specified		A 100 M 100
. 70	A THE RESIDENCE SHAPE	Philadelphia and the second	1	his average would be 13 015		100000000000000000000000000000000000000

# QUARTERLY METEOROLOGICAL TABLE,

Compiled from the Weekly Tables furnished to the Registrar General by the Astronomer Royal.

TABLE,

Deaths registered in the Metropolis from Small Pox, Measles, Scarlatina,
Hooping Cough, Typhus, Diarrhea, Dysentery and Cholera, Influenza,
Consumption, and other diseases of the Lungs; the numbers at each
of the Astronomer Royal.

by the Astronomer Royal.

the 13 weeks ending March 30, 1844.

Highest, of 13 weeks.	30th	23rd	16th	9th	h 2nd	24th	17th	10th	ary 3rd	2716	20th	13th	ry 6th	cek	© 44 #4
	28th,	19th,	11ић,	4th,	Feb. 26th,	18th,	11th,	6th,	Jan. 31st, deci	27th,	19th,	12th,	5th,	> 0 H	
SEUL	full	new	3rd qr.	full	1st gr.	new	3rd qr.	In equator	Greatest lination N.	1st qr.	new	3rd qr.	full	Phases of the Moon	
29.705	29.948	29.704	29.544	29.798	29.237	29.363	29.951	29.269	29.706	30.042	30.059	30.068	29.479	Mean height of the Baron 72 observations, correcte duced to 32 deg. Fahrenh	eter, from
57.6	57.6	49.4	071 823 64	55,6	51.1	48.3	47.0	43.0	50.1	47.4	46.5	43.8	52.8	Highest during the week	
6 18.6	33.7	29.0	32.6	24.3	28.0	23.5	21.2	25.8	27.5	30.2	26.3	32.9	18.6	Lowest during the week	
49.7	54.6	46.5	46.5	42.8	45.9	41.7	40.1	39.4	40.5	43.2	40.3	41.5	45.2	Highest on each day, 6 observations	
39.4	40.8	33.0	35,9	31.9	34.6	29.1	28.7	30.7	33.0	34.2	33.2	35.1	33.7	Lowest on each day, 6 observations	Mean
10.4	13.8	13.5	10.6	10.9	11.3	12.6	11.4	8.7	7.5	9.0	7.1	6.4	11.5	Difference	1 10 10
38,6	47.2	39.7	41.1	37.3	40.3	36.5	34.7	34.6	36.9	39.2 3	37.0	38.8	40.1	Mean of 72 observations	- 0
34,9 8	42.5	33.7 6	36.7 67	32.7 6	35.6	31.6 57	32.1	32.2	32,5	36,3	33,3	36.6 51	37.9 5	Mean of 72 results	Dew Point -
83.6	83.6 7	62.2 5	0	68.7 5	69.0 6	57.2 4	56.0 40	58.0 41	61.4 4	52.3 4	49,0 4	51.0 4	53.6 46	Mean of 7 observations	1 2
52.3	70.1 26.	58.3 16	57.2 24	58.4 15	60.7 22.1	47.7 12.	46.3 14	48.6 17	49.2 18	47.9 21	43.6 15	45.7 24	46.5	During the week	Self-Registering
6.3 27.1	33.4	16.5 27.1	24.0 29.9	15.2 26.4	29.3	2.8 24.4	14.0 23.1	17.2 24.3	18.6 25.6	21.2 28.4	15.2 24.8	24.2 29.5	6.3 26.2	Mean of 7 obser-	tering
	-	-	9							-				vations • • • • • • • • • • • • • • • • • • •	
	4.8	6.0 10	.4 9	4.7 9	4.7 10.7	3,9 9	2.6	2.4 6	- A	2.0	3.8	2.3	2.2		Difference be- tween the dew point temperature and air
7.0	10.3	10.7	9.2	9.6	17	9.4	6.0	6.2	8.0	5.7	6.8	sia.	S. S.	Mean of the greatest on each day, 6 obser- vations  Mean of the least on	erence n the d temper and air
0.8	8.0	1.6	0.5	1,2	1.4	0.6	0.7	1.0	1.3	0.8	1.0	0.5	0.2	Mean of the least on each day, 6 observa- tions	lew ature
2.	+4.6	12.8	111	-3,5	-0.0	-4.0	-4.1	-4.6	-0.2	+1.7	+2.1	+3.1	+ 4.5	Difference between the mes ature of the week, and temperature of the same an average of 25 years	n temper the mea
	s.w.	N.W.	N.W.	3	w.	Variable	N.W. and S.W.	W.S.W.	W.N.W.	s.w.	N. & W.	W-S.W.	s.w.	General direction	Pressure
17.0	7.0	8.0	17.0	9.0	10.0	19.0	1.0	4.5	12.0	1.5	C0 67	1.5	8.0	Greatest pressure in the week	bs. D
0.69	0.5	1.0	1.7	0.9	1.2	0.5	0.1	0.2	1-2	0.1	0.3	0.07	0.4	Mean for the week	on the
1238	1180	1385	1660	1440	1860	1460	750	970	1345	610	865	930	miles 1645	The amount of Horizontal of the air in each week	m <b>ovem</b> en
7.2	6,4	7.6	7.3	6.9	7.2	7.0	8.1	6.0	6.7	7.5	7.5	8.6	7.0	Mean amount of Cloud, 0-10	
7-13	0.04	0.13	0.97	1.25	0.69	0.73	0.02	0.60	0.40	3	3	1.06	1.24	Rain in inches (7 days)	
	18	22	21	16	17	19	29	17	19	18	21	20	16	Small Pox	70112
	25	27	10	. 32	29	80	100	30	28	23	26	23	25:	Measles	
	83	36	43	40	39	40	43	37	31	42	40	51	47	Scarlatina	
	4	51	83	49	52	33	37	44	97	89 80 80	68	82	228	Hooping Cough	
	39	19	28	. 91	41	42	40	30	38	33	35	34	23	Typhus	
	13	7	0	0	13	00	10	4	6	80	12	6	10	Diarrhota, Dysentery, and C	holera
Maria Maria	9	4	w	7	۵.	*	9	Cn Cn	-	4	ω.	w	CO	Influenza	
	140	147	193	150	156	166	170	166	162	132	131	130	111	Phthisis or Consumption	
	152	162	174	225	210	263	278	249	229	214	217	193	174	Other diseases of the lungs.	
	44 63 64	427	398	443	450	410	463	452	444	432	450	431	440	0 to 15	Age
	310	304	.930	.326	336	392	361	350	358	322	354	287	259	15 to 60	s, (excl of and Deaths
	205	209	229	238	236	293	\$82	243	246	235	229	193	190	60 and upwards	Ages, (exclusive of violent and sudden Deaths)
	958	948	957	1008	1024	1087	1108	1045	1052	166	1013	911	903	Deaths from all causes, (exviolent and sudden deaths)	

The Ages of 39, on account of the corrections for returns not received,
were not specified in the Table.
 Inquests are held on violent and sudden deaths which are returned irregularly.

Mean, H or Lov the 13

### REMARKS.

The District of Wandsworth was added to the Metropolis at the beginning of the Year 1844; and the Return for the present Quarter is derived from 115 Districts, divided into 576 Registrars' Districts. Thirty-four of the Districts are included in the Metropolis; and the remaining eighty-one Districts comprise, with some Agricultural Parishes, the principal Towns and Cities of England. The Population was 6,578,912 in 1841.

Table I.—46,041 Deaths were registered in the Winter Quarter of 1844; namely, 13,471 in the Metropolis, and 32,570 in the other eighty-one Districts; the average of the corresponding Quarters of 1838-42 having been 13,266 for the Metropolis, 32,085 for the other Districts, 45,351 for the Metropolis and the other Districts together. The Deaths were, therefore, 690 above the average of those five Winters. 400 of the excess being substracted for the additional day in Leap-year, 290 are left.

But the Population has been found at all the Censuses to be increasing rapidly in the Town Districts, partly by the excess of Births over Deaths, and partly by immigration. The Population of the fifty-nine most populous Districts in the Return increased at the rate of 6.3 per cent. every four years, from 1831 to 1841; and it is, therefore, probable that the Population did not increase less than 5 per cent. in the four years from the Winter Quarter of 1840 to the Winter Quarter of 1844. The average with which the 46,041 (—400) Deaths in the Quarter ending March 31st, 1844, should consequently be compared is 47,619.

Table III.—The Metropolis has been less healthy than in previous years; but the mortality is not so much above the average as in the Autumn Quarter of 1843. The average employed in the Table was derived from the Returns of the five years, 1839—43. It will be observed that 252 persons died of Small-pox, which was fatal to only 114 persons in the preceding Autumn Quarter. The great majority of these persons were Children, and had not been vaccinated. Measles and Scarlatina were declining, yet 334 persons died of the former, 536 of the latter disease. Influenza, Bronchitis, and Pneumonia, destroyed more than the average number of lives in the Metropolis; and their fatality, particularly among the aged, is mentioned in many of the Reports from the other Districts. The Deaths registered from Dropsy, Cancer, Scrofula, Heart Disease, Apoplexy, Paralysis, Tabes Mesenterica, were above the average in number; those from Childbirth, as well as violent and sudden deaths, were below the average. Two deaths happened from Hydrophobia.

The Quarterly Meteorological Table has been compiled from the valuable weekly Tables supplied by the Astronomer Royal. Upon comparing the results with those obtained from the observations of Mr. Henry, it will be seen that the mean temperature of the air has been a little below that of the season.

Districts in which the Mortality was higher than the average Mortality of Winter in the same Districts.—Brighton, Winchester, Windsor, St. Albans, Wycombe, Northampton, Bedford, Cambridge, Yarmouth, Dorchester, Exeter, Plymouth, Kidderminster, Birmingham, Aston, Coventry, Lincoln, Wigan, Abergavenny, Merthyr-Tydfil, Anglesey.

Districts in which the Mortality was lower than the average Mortality of Winter in the same Districts.—The Central Districts of the Metropolis, Cheltenham, Walsall, Wolverhampton, Stock-port, Macclesfield, Preston, Bolton, Prescott, Manchester, Salford, Sheffield, Leeds, Sunderland, Gateshead, Newcastle-on-Tyne, Cockermouth, Pontypool,

# REGISTRARS' NOTES TO THE MARCH QUARTERLY RETURN.

The following Direction was addressed to the 447 Registrars in the Country:—

"If at any time the number of deaths registered during the Quarter has been above the average, state, in a note at the foot of the Return, whether any epidemic disease, such as Measles, Typhus, &c., has been prevailing in the District; or if there be any other known circumstance which will account for the increase."

In compliance with the above direction, 206 Registrars have appended explanatory notes to their Returns; the remaining 241 Registrars apparently observed a decrease, or no obvious increase, in the mortality.

An Abstract of the Notes to the Returns is subjoined.

REG. DISTRICT.

SUP. REG. DISTRICT.

MAIDSTONE Loose	The number of deaths this quarter is above the average, which may be attributed to the prevalence of <i>Hooping Cough</i> and <i>Inflammation of the Lungs</i> among children, and a considerable number are from old age.
BRIGHTON The Palace	An increase upon the two preceding quarters, although no particular disease has been prevalent. There have been 17 deaths from Consumption; 25 persons were at advanced ages—6 whose united ages averaged 53 years each—6, 66 years each—8, 75 years each—3, 84 years each—1, 93—and 1, 96 years.
Kemp Town	The return is 23 above the average of five years, which I attribute to a greater number of visitors than usual, there being no prevailing disease.
PORTSEA ISLAND Portsmouth Town	vaccination—and 2 arter vaccination, cc.
Porisea Town	The deaths exceed those of the corresponding quarter last year by 12. Of the deaths registered this quarter, 10 are from Small Pox—one after, and nine without previous vaccination. Twenty-nine of the number registered were children under 3 years of age, and 20 of them adults from 60 to 86 years.
Kingston & Landpor	The mortality compared with corresponding quarters is below an average, although the Government contract works and the building the new union house, have caused a considerable increase in the population. The immates of the workhouse have been very healthy. There have been but few cases of epidemical disease, and, I believe, at the present time, my district may be said to be very healthy.
WINCHESTER Winchester	The state of the s
WINDSOR Windsor	Rather above the average, the increase arising chiefly from deaths by old age, &c. Of 82 deaths, 27 were of persons above the age of 60—19 above 70—8 above 80—and 2 above 90
Egham	Above the average-16 more than the corresponding quarter of 1843, but no particular disease has been prevalent.
ST. ALBANS St. Albans	The number of deaths in my district is 20 more than in the corresponding quarter last year. This has arisen from the deaths of an unusual number of aged persons. The united ages of twelve females amount to 907 years.
WYCOMBE High Wycombe	The number of deaths is 15 more than in the corresponding quarter of last year. It includes 5 coroners' inquests, but no particular cause can be assigned for the remainder, as I am not aware that any epidemic is prevalent in this district. They are of very old persons and infants.
Great Marlow	The number of deaths this quarter is less by 10 than the last quarter. Measles is less prevalent in the town of Great Marlow, only 1 case having proved fatal. The disease is extending to the agricultural part of my district, 6 cases having been fatal. I have registered 13 cases of old age from 60 to 90 years; the other cases are of the ordinary description.
OXFORD Oxford	There is no particular cause to be assigned for the 6 over the average.
NORTHAMPTON All Saints	There has not been so great a number of deaths in my district in any one quarter since March 1835. The average of the 7 corresponding quarters is 70½, including the present one of 83 deaths. Measles has been, and still is, very prevalent. Births this quarter 120.
St. Giles	There is still an increase in the number of deaths; this quarter Measles and Influenza have been prevailing in the district, and in a few cases proved fatal. The increase may not be considered alarming, having the Union Workhouse, Lunatic Asylum, General Infirmary, and Barracks situated in the district.
BEDFORD Bedford & Carding	ton Number of deaths below the average. The district generally very healthy.
	on The number of deaths registered this quarter is 2-11ths above the average number registered in the corresponding quarters of the preceding six years; but this increase of mortality is not attributable to any epidemic. Influenza has been prevalent but not fatal,
CAMBRIDGE St. Andrew the Le	ess The number of deaths this quarter, as well as the last, is double the average. The chief cause is Small Pox, from neglect of vaccination. In 30 deaths only 2 doubtful cases have occurred after vaccination.
St. Andrew the Gre	at The number of deaths is considerably above the average. Small Pcx, Measles, and Inflammation of the Lungs have been fatal in many cases.
St. Giles	There is considerably more than double the amount of deaths registered this quarter than in the corresponding one of last year. Small Pox and Fever have been prevalent in this district. Out of 15 cases of Small Pox only 1 was after vaccination.

		The state of the s
SUP. REG. DISTRICT  IPSWICH	REG. DISTRICT. St. Matthew	The number of deaths is above the average. I have registered more deaths from old age than in any preceding quarter. Hooping Cough has been prevalent and fatal.
NORWICH	Conisford	The number of deaths this quarter has been above the average. No epidemic disease has been prevailing in the district, but the causes of death of 17 persons have been registered as being Old Age and 10 Phthisis.
********	Coslany	Above the average number, but all from the generality of causes.
GREAT YARMOUTH	Northern	The number of deaths registered this quarter is 18 above the corresponding quarter of 1843. There are 41 cases of Affection of the Langs; Bronchitis has proved fatal to children and old people; there are 17 cases of Phthisis; of Scarlatina there are 16 cases, and of Typhus 7. The number of deaths at 5 years and under is 43; of 60 years and above, 33. Deaths are recorded of persons who have reached the ages of 80, 81, 82, 84, 86, 90, 94, and 95 years.
2.3 % - (6.56 · )	. Southern	The number of deaths registered this quarter is considerably above the average. Many children have died from <i>Pneumonia</i> and <i>Scarlatina</i> , and 21 persons from the age of 70 to 90.
DEVIZES	Lavington	15 deaths less this quarter than the last, in consequence of Scarlatina and Typhus being less prevalent.
		Rather above the average; out of the above number, 2 were of Scarlatina, 4 of Typhus, and 3 Coroners' Inquests.
DORCHESTER	The state of the last	The number of deaths this quarter exceeds the average; the increase is mainly attributable to Scarlatina at the commencement of the quarter.
EXETER	St. David	Above the average, but from no prevalent disease. The mortality among children noticed in the last quarter's return continued during the month of January.
	St. Sidwell	Above the average; attributable to Pulmonary Inflammations, both idiopathic and accompanying Hooping Cough.
ST. THOMAS	St. Thomas	Among the number were 15 (principally females) of the ages following—I of 73; 2 of 74; 3 of 80; 1 of 81; 2 of 82; 3 of 85; 1 of 86; 1 of 87; and 1 of 95 years. 2 cases of Measles, 1 of Hooping Cough, and 4 inquests.  Seven of the deaths were from Hooping Cough, which has been and is still very
***************************************	Kenton	prevalent. There have been two Coroner's Inquests.
	Heavitree	This return is above the average; several persons of advanced age have died. In almost all of the cases it was stated to be from natural decay. Influenza is very general, and 4 have died from Hooping Cough. Only 1 case of Typhus registered and 4 inquests, the remainder Chronic diseases.
The state of the state of	Alphington	The number of deaths is 12 more than in the corresponding quarter of last year, and exceeds by 11 that of the last quarter, attributable to the prevalence of <i>Hooping Cough</i> and <i>Measles</i> . This return includes 6 cases of <i>Hooping Cough</i> , 7 of <i>Consumption</i> , and 2 Coroner's Inquests.
20 - 12 - 12 - 13 - 13 - 14 - 14 - 14 - 14 - 14 - 14	Woodbury	There are two deaths less than in the corresponding quarter last year. The district very healthy, no epidemic prevailing.
PLYMOUTH	. Saint Andrew	The epidemic of Small Pox has tended to raise the mortality above the average of the winter quarters, 25 deaths are attributed to that cause; there have been also 5 deaths from Measles, and more than the average number from diseases of the respiratory organs, as well as from old age or diseases incident thereto.
di 24 (2001-0	Charles the Martyr	The deaths are below the average. Small Pox has prevailed considerably: in almost all the fatal cases the sufferers were unvaccinated.
REDRUTH	Redruth	Eighteen of the 53 deaths have been produced by Consumption. Two cases of Typhus and 2 of Scarlatina only.
	. Phillack	Ten above the average, owing to 10 deaths from old age, and 12 from Pulmonary Consumption.
PENZANCE	St. Buryan	My district is tolerably healthy at present; there is a decrease as compared with the corresponding quarter of last year of 7.
	. Marazion	The return is above the average, owing to the prevalence of Scarlatina, which is still in the district.
BATH	Lansdown	The number is 9 above that of the corresponding quarter of 1843; 5 of the deaths are from Scarlatina, and 5 from Hooping Cough.
	Walcot	The return this quarter is above the average, owing chiefly to mortality amongst old people. There were some few fatal cases of <i>Scarlatina Maligna</i> , and <i>Hooping Cough</i> the early part of the quarter, but at present we are perfectly free from any epidemic.
***********	. The Abbey	The return is above the average owing to the prevalence of Scarlatina.
, , , , , , , , , , , , , , , , , , , ,	· Lyncombe	Scarlatina has prevailed to a great extent, principally of the miliary kind, accompanied with enlargement of the parotid glands; several cases have terminated fatally.  Anasarca in some cases has followed the Rash, with affections of the joints; Measles has been prevalent, and Hooping Cough has followed in several cases. I have seen more of the above cases this last quarter than for three or four years preceding.
	. Twerton	The number of deaths in my district for the quarter ending March 31st is above the average: I have registered 10 cases of Consumption, 3 of Scarlatina, 6 of old age, and 5 of Typhus.
BRISTOL	. St. Augustine	A greater proportion than usual of old persons, 20 being 60 years of age and upwards, principally from the gradual decay of nature, with Asthma, &c. One half the whole number are children under 7 years of age, a less proportion than usual. Scarlatina prevalent—fatal in 7 cases only.
	. S James	The number of deaths is 12 less in this quarter than in the last, but 15 above the average of last year. This has been principally owing to 39 cases of Diseases of the Lungs, Phthisis, Pneumonia, &c. many of which were sent to the Infirmary, not only from different districts of the city, but probably from the neighbouring parishes.
	. St. Paul	Scarlatina continued to prevail in the early part of the quarter has not exceeded the average.  Scarlatina continued to prevail in the early part of the quarter; 6 fatal cases have been registered in this district. Small Pox also has appeared to some extent, and continues in one locality; two fatal cases (without previous vaccination) have been registered.  Many cases, however, have presented themselves in which vaccination had been previously performed; but in all these the disease appeared to be modified and exceedingly mild in its character.
CLIFTON	. St. Philip and Jacob	
	. Ashley	The last quarter has been most unhealthy. The number of deaths is much above the average. Out of the amount registered there are 4 from Scarlatina, 2 Small Pox, 2 Typhus, and 2 Measles.
		the state of the s

SUP. REG. DISTRICT	REG. DISTRICT.	Action of the Control
CLUFTON	. St. George	There has been a great amount of mortality this quarter in my district; and if the unusual number of deaths that occurred at the Union Poor-House for Children in the quarter ending Sepfember 30th, 1842, and the 1 death that occurred there this quarter, be both taken out of the account, the mortality in all the other resident families this quarter will greatly exceed that of any previous quarter; taking the 68 deaths registered and 2 Inquests (one from natural cause) not yet registered, the increase this quarter is 58 per cent. above the general quarterly average, being at the annual rate of 3.25 per cent. in the population, estimated to be now 8600. The increase has been caused by the following epidemics:—the deaths being from Scarlatatina 15, Typhus 3, Diarrhea 2, Measles, 3, and Hooping Cough 2. There have been 7 deaths from Consumption, and 13 children, not included in any of the before-mentioned cases, have died under a month old.
STROUD	. Stroud	The increase of the number of deaths this quarter has been in a great measure amongst those advanced in years. Hooping Cough, Measles, Scarlatina, Inflammation of the Throat, and Rheumatism, have prevailed considerably in the District, with a few cases of Small Pox, rarely fatal.
.,	. Painswick	The number of deaths is above the average this quarter; a third of the deaths registered were caused by <i>Pulmonary Consumption</i> .
HEREFORD		The increase of deaths appears to arise from the death of children under 5 years of age.  The deaths this quarter are below the average, although Measles has prevailed very much in the district, and has been fatal in a few cases.
	- Madley	Rather above the average of the winter quarter of 1843. Influenza has been very prevalent, and in some cases fatal.
SHREWSBURY		Although 20 above the average, and 25 more than last quarter; still there are few cases of Epidemic or Contagious Diseases. 3 of Measles (first appearance for a considerable time in my district), I Hooping Cough, I Small Pox (without previous vaccination). Consumption (12 cases) forms a considerable portion of each quarter's return, being above a tenth of this quarter, Convulsions amongst children (9) forming nearly another tenth part of the whole return. Total 100 cases.
KIDDERMINSTER .		It appears to me that the great increase in the number of deaths is attributable to Consumption and Inflammation of the Bowels. The former disease has proved fatal in 32 out of 159 deaths, which is 1-5th of the total number for the quarter.
The same of the sa	Lower Mitton	
DUDLEY	. Rowley Regis	Influenza and Hooping Cough. These diseases have been very prevalent in this district. The number of deaths this quarter is nearly double the amount registered in the corresponding quarter of last year, arising chiefly from the fatal prevalence of Typhus and other kinds of Fever; last quarter 42 deaths were registered, being about the average for 1843.
WALSALL	The Borough	There is an increase of 33 deaths this quarter. 19 have died of Consumption, and 22 of Inflammation; these two complaints have been most prevalent.
- 10 Table 1	Darlaston	The number of deaths in my district rather exceed the usual average this last quarter;  Inflammation in the Bowels is the most prevailing disease.
WOLVERHAMPTON		The deaths this quarter are 43 more than in the corresponding quarter of 1843. There has been no epidemic. The increase is owing to a greater number of deaths amongst persons advanced in life, of Asthma and Chronic Diseases. There is one peculiarity of this quarter, that is, the excess of males over females, being 22.
	Willenhall	The number of deaths is above the average. This quarter 9 have died of Fever, 10 of Consumption, 8 with Inflammation in the Bowels, 12 Inflammation of the Lungs or Chest.  Seven more than the average. Influenza has been prevalent during the last
eds all a margaret att	and a state to the same of	quarter, and has generally proved fatal in its attacks upon old people. 5 deaths out of 24 from Influenza in the above district during the last quarter.
WOLSTANTON	Burslem	The number of deaths this quarter is above the average of the year, and is also above the number of the corresponding quarter of last year to the extent of 5, there having been in this quarter 8 more deaths of males and 3 less of females than in the March quarter 1843. No epidemic has prevailed to any extent in the district; 4 deaths have been caused by fever, 1 from Typhus, 1 from Brain Fever, 1 from Scarlatina, and 3 from diseases consequent upon childbirth. The principal causes of death have been Diseases of the Chest and Lungs.
BIRMINGHAM	St. George	During the quarter in my district 11 have died of Typhus, 23 of Scarlatina, 24 of Consumption, 36 Inflammation of Chest and Lungs, 11 Inflammation of the Bowels, 15 Hooping Cough, 23 of causes investigated at Coroners' Inquests, the remaining 98 from sundry other causes.
	St. Paul	There have been this quarter 32 deaths above the average in this district; the prevailing diseases were Consumption and Inflammation of the Chest, Lungs, and Bowels. Consumption 12, Inflammations 22, Scarlatina 3, Typhus 3, Hooping Cough 3, Measles 2, Influenza 1, Bronchitis 2.
	Lady Wood	There has been in this district an increase of deaths this quarter over former quarters, including considerable increase of epidemics, especially <i>Fever</i> , without any particular known cause, unless mildness of the winter may be supposed such.
ASTON	Duddeston	The number of deaths this quarter is above an average; Measles, Croup, and Hooping Cough have been prevalent.
	. In ungerthe	The number of deaths in this quarter, January to March 1844, has been above the average as compared with the last two years, to which only I can refer, the number for 1842 being 343; number in quarter ending March 31, 47; 1843; 33; number in quarter ending March 31, 30. 11 persons died under 2 years of age, 19 under 60, 3 of Dysentery, 3 of Scarlatina, 2 of Typhus, 3 of Hooping Cough.
	Deritend	The number of deaths registered this quarter is rather above the average, which I account for by observing that Scarlatina has been unusually fatal to children (10), and Inflammation of the Lungs to aged persons. 23 of the number registered were persons above 50 years of age.
COVENTRY	Holy Trinity	The above is rather a higher rate of mortality than is usual in my district. Influenza has been particularly prevalent, and, as I am informed by medical men, has operated, to a great extent, as the cause of disease terminating fatally.
LEICESTER	East Leicester	There is an increase of 31 deaths over the corresponding quarter of 1843, principally amongst old people, and Inflammation of the Respiratory Organs. I consider the district healthful, no epidemic prevailing.

Mortality (123) in Lincoln has been considerably on the increase. My return of deaths for the quarter ending September 1842 was 80—each succeeding quarter has shewn an increase; the past quarter's return exceeds the previous returns by a fifth. This excess of mortality is not attributable to any prevailing epidemic. Fever, though exceedingly prevalent, has not been very fatal. The mortality among infants, accounted for by the general cause—"Convulsions"—has been considerable, numbering 34 under 1 year of age; the great inortality has been among the vigorous in years. In 22 cases Consumption has been fatal, and in most instances in the prime of life. Lincoln has been and is celebrated for its salubrious atmosphere. The general good health of its inhabitants has been proverbial. Whence then, it will be asked, is the cause of the increased mortality? In a report 1 had the honor to make some time back, I alluded to the great want of sewerage—creating large and numerous deposits of vegetable and animal matter in a state of decomposition, producing miasma and its certain consequence—continued fever. It is a curious fact, that the intermittent fevers in former years so prevalent in these districts, as to render Lincolnshire ague a proverbial expression, have, under the influence of steam-drainage of the fan lands, disappeared almost entirely, while the NECLECT OF TOWN SEWERAGE, above noticed, has increased the prevalence of what is here called continued, or Typhus fever, to an extraordinary amount. At the annual meeting of the Dispensary held this day the following view of a decennial period appears in the report. Their year commences and ends on the 25th of March. SUP. REG. DISTRICT. LINCOLN ...... Home ... Cases treated of Continued or Typhus Fever, and Intermittent. 1836 1837 1838 1839 1843 1844 1840 1841 1842 Intermittent ..... 15 7 7 5 4 0 Lincoln has every capability of appliance for carrying out the most complete sewerage—its necessity is admitted by all, yet it requires a Legislative enactment to do that which common prudence suggests, and which the natural protection of human life The average number of deaths is  $55\frac{1}{2}$ . The deaths registered this quarter are Typhus 6, Hooping Cough 11, Measles 5, Influenza 3, Consumption 8, other causes 46, Total 79. South ...... The number of deaths this quarter is 65, being 11 less than the average. NOTTINGHAM .... Byron ..... Ten of the deaths have been from measles. .... Saint Mary ...... Rather above the average, but no epidemic has prevailed, 1 case of Typhus and 1 of Scarlatina being the whole of that nature of disease registered during the quarter. BASFORD ..... Arnold ..... The return presents a considerable increase upon several previous quarters, arising chiefly from the deaths of aged persons; the district is generally in a healthy state. Bulwell ..... In the return for this quarter it will be observed, that the number of deaths among males has greatly exceeded that among females—nearly in the proportion of 2 to 1. The mortality in children under 2 years of age has been very remarkable, the majority having been carried off by convulsions. The number of registered deaths last quarter was 59, so that the present number (80) shows an increase of 21. Typhus and Scarlatina have both been prevalent, but the number been principally augmented by Phthisis and diseases incidental to age. Basford ..... DERBY ...... St. Peter ...... The number of deaths in this quarter is 34 more than in the corresponding quarter of has last year, and 8 less than in the same quarter of the year 1842. STOCKPORT ...... Hyde ...... District generally healthy. Although this has been a sickly quarter the deaths show a decrease of 22 upon the preceding one, which may be attributed to the total absence of Scarlatina. Influenza has been very prevalent, but of so mild a form as not to cause I fatal case; with this exception I am glad to report my district free from epidemics. It may not, perhaps, be uninteresting to compare the deaths from Consumption for the last four quarters; they are, consecutively, 19, 18, 10, 18. The total deaths are 375; so that the mortality alone from this "fell destroyer of the human race," is 17 1-3rd per cent. or rather more than one death to every sixth. Measles 1, Diphtheria 1, Hooping Couph 2, Thrush 1, Croup 1, Diarrhea 1, Typhus T, Dropsy 6, Consumption 18. Of the 86, 55 were attended by surgeons, 5 by the Infirmary, 12 by druggists, 10 had no medical aid; the remaining 4 were Coroners' Inquests. .... Cheadle ..... • Below the average. For the quarter ending March 1842, I registered 26 deaths, ditto 1843, 31 ditto, 1844 only 14. - ...... Hazelgrove......... In this quarter there are 2 less than in the last quarter, and 6 less than in the corresponding quarter of 1843. The district is very healthy at present. The last month there were many cases of Pleurisy, but none proved fatal. MACCLESFIELD .... Macclesfield East ..... The number of deaths this quarter is 115, being a little above the average. The prevailing diseases are Epidemic Influenza, Consumption, Inflammation of the Lungs, or Pneumonia and Fever. The number of deaths in the present quarter is upwards of 20 per cent. above the average. There has been no particular epidemic prevalent. The increase has been generally owing to the common causes of mortality. .... Sutton ..... The number of deaths this quarter is above an average, being 20, but not from any epidemic disease. 3 old people upwards of 80 years, 6 upwards of 60 years of age, some of them with a very short sickness, 2 from Teething, Consumption, &c. · ... Alderley ...... The deaths in my district this quarter are 57, being 12 above the average. The greatest number of deaths occurred in the month of February; the increase is chiefly among the aged, 15 having died above the age of 55; viz. 3 between 55 and 60; 3 between 60 and 70, 5 from 70 to 80; and four at the ages of 80, 82, 86, 90. GREAT BOUGHTON. Hawarden ..... -. The Castle .......... 28 deaths registered more than in corresponding quarter of last year; and exactly that number more on the average of the 4 quarters ending December 1843. The increase is chiefly confined to infants, consumptive, and aged persons.

SUP. REG. DISTRICT. REG. DISTR	ICT
	reet The return is under the last quarter. It includes 21 cases of Small Pox, 3 only of which were vaccinated. They were all children of the lower Irish, who have not a very
Dale Street	distinct idea of the nature and benefits of vaccination, and lamentably neglect it.  The deaths in my district have been very regular, only 6 more than the previous
St. Thomas	quarter, the increase may be ascribed to Consumption.  There is a decrease of 10 on the quarter 1842, and an increase of 20 on the quarter 1843.  The following are the principal complaints of the quarter, 158 of which were under 5 years of age:—Pneumonia 32, Phthisis 35, Bronchitis 20, Hooping Cough 21, Convulsions 34, Fever 16, Measles 6, Small Pox 2, Inquests 13, Croup 5, Diarrhoea and
Mount Pleasant .	Enteritis 8.  The number of deaths in my district this quarter is 63 more than the last, and 57 above the corresponding quarter of 1843; the most prevalent causes of death appear to be Inflammation of the Lungs, Dropsy, Fever, Dysentery, Asthma, Paralysis, Consumption, Disease of the Heart, Measles, Bronchitis, Apoplexy, Marasmus, and Hooping Cough.
Islington	There are 68 deaths less this quarter than in the corresponding quarter of 1843. I have registered 44 deaths from Phthisis, 26 from Pneumonia, and 20 from Hydrocephalus.
Saint George	increase I attribute to the prevalence of diseases of the Heart, Asthma, and Phthisis in aged persons.
WEST DERBY West Derby	There is a decrease of 43 in the number of deaths, as compared with my last quarterly return. Particulars:—Decline 5, Convulsions 11, Dropsy 5, Water on the Brain 6, Consumption 7, Decay of Nature 3, Palpitation of the Heart 3, Putrid Sore Throat 1, Scarlet Fever 3, Inflammation of the Lungs 11, Fever with Aboess 1, Dysentery 1, Childbirth 1, Asthma 4, Aneurism of the Aorta 1, Hypertrophy of the Heart 1, Diarrhosa 1, Inflammation of the Bowels 1, Tubercular Consumption 2, Croup 1, Liver Complaint 1, Cancer 1, Apoplexy 1, Phthisis Pulmonalis 2, Premature Birth 2, Small Pox 1, Teething 1, Brain Fever 1, Inquests 4.
	The return presents a decrease of 6 deaths compared with the last quarter, though Typhus has been raging in Ince Blundell, occasioned, it is presumed, by lodgments of filth in various parts of that township.
Toxteth Park	the mortality being the same as in December quarter 1843.
BLACKBURN Blackburn	The deaths this quarter are about the average number, but less by 130 than the same quarter last year. We have had no particular epidemics. Only a few cases of Measles and Hooping Cough. 60 infants have died in fits, and 66 persons of Consumption.
Darwen	••••• There has not been any spidemic disease in my district during the past quarter. We have had 1 death more than in the corresponding quarter of last year.
Witton	The number of deaths in this quarter is the same as the last, which shows that my district is in a healthy state, taking into consideration the season of the year.
	There are 5 deaths less this quarter than the corresponding quarter; the district healthy The number (48) is 2 more than the March quarter of 1843; but is seven less than the
ROCHDALE Whitworth	average of the March quarters since the commencement in 1837  The number of deaths in my district is above the average of former quarters, owing to
Wardleworth	average, I having had no less than 21 cases of Small Pox, without previous vaccination,
Weurdle	and not one that had been previously vaccinated.  This quarter's return is rather above an average, caused by Measles and Convulsions having been prevalent among children; likewise more cases of Consumption than usual.
BURY Bury, North	***** The deaths occurring in my district are considerably above an average. Measles and Scarlatina have prevailed.
—— South	The return shows an increase of 27 in the number of deaths registered during the present quarter over those registered in the corresponding quarters in the years 1838, 1839, 1840, 1841, and 1842; the whole number registered during the present quarter being 117. Measles have been very prevalent during the whole quarter, from which disease there has been 43 deaths, 20 males and 23 females; and from Scarlatina 7, males 4, females 3; there have also been 6 Inquests, 5 over persons who have died suddenly from natural causes, and 1 from accident.
	There has not been any particular epidemic, but a number of deaths more than in the quarter ending December 1843.
Elton	····· Above an average number; there have been 10 fatal cases of Measles, which account for the increase.
BOLTON Eastern	Typhus and Hooping Cough appear to have prevailed, so as to cause a small increase above the average.
Western	The number of deaths in my district is above the average of past quarters, owing, I believe, to the increase of population (as many houses which were uninhabited are now inhabited) and not to any epidemic which has prevailed. Births also have increased in proportion.
	This quarter is about 1-4th more than the average; there have been several cases of Croup, which appear to have been the cause of increase.
Edgeworth Farnworth	····· Measles and Hooping Cough have been prevailing in this district.
Tong with Haulg	The return of deaths is below the average of corresponding quarters of former recess
	average, from which may be inferred that my district has been in a very healthy condition during the past winter.
WIGAN Wigan	294 have been registered, and out of this number 181 have died under 5 years of age; 11 from Measles, 12 from Croup, 30 from Scarlatina, and 59 from Small Pox, without previous vaccination.
Standish	****** 49; the deaths for the same quarter in 1843 were 32; thus there is an increase of 17. the prevailing diseases have been Debility, Scarlatina, Old Age, Convulsion Fits, Small Pox, and Consumption.
Aspull	•••••• Number of deaths this quarter 57, an increase of 27; Typhus and Scarlatina 7, Measles 3, Teething 6, Inquests 5, Fits and Debility 11; Old Age 7, Chin Cough 1, Croup 2; upwards of 33 children having died principally under 2 years of age, of diseases incident to them.

SUP. REG. DISTRICT.	REG. DISTRICT.								
	Hindley	There has been more than an average number of deaths in my district for the last quarter on account of Measles and Small Pox having been prevalent.							
********	Ashton	The deaths registered in my district exceed the average of the last 2 years, owing to the prevalence of Pneumonitis, Bronchitis, Croup, and Small Pox.							
	Upholland	10 per cent. above the general quarterly average. No epidemic has been prevalent; but some cases of Small Pox have occurred in a part of the district, terminating fatally in the ratio of 10 per cent. upon the number registered. The deaths of children under 5 years old are 53 per cent., and of persons above 60 years old 21 per cent.							
PRESCOT		Prevailing epidemic Influenza. Deaths equal to the corresponding quarter.							
*********		This district has been very healthy this quarter.							
	Prescot ,	The number of deaths registered in this quarter, ending 31st March, 1844, is 89, being about 27 above the general average; there are 21 entries of deaths under 2 years. The sudden changes in the weather have been very fatal to old persons; there are 17 entries of persons who have died between the ages of 50 and 70 years, 7 entries between 70 and 80 years, and 4 above 80 years.							
£+0+;+++	Huyton	My district is below the average of last quarter.							
	Much Woolton	The number of deaths registered in this district during the present quarter is exactly the same as in the first quarter of the operation of the Registration Act, both being more than any of the intermediate quarters. The increase in the first was caused by the prevalence of Small Pox, and in the last by the deaths of aged people. The united ages of 9 females amount to 640 years, and of 4 males to 240 years. There have been some cases of Small Pox in the district, but only I has proved fatal.							
	Hale	In the last 4 quarterly returns the deaths are much below the usual average. In the township of Hale, containing, by the last census, a population of 645 persons, not one death occurs between April 26th, 1843, and January 30th, 1844, a period of more than 9 months.							
CHORLTON	Chorlton on Medlock	Though there has been, I believe, rather more sickness than usual during the past quarter in this district, the mortality has been somewhat below the average amount. This is no doubt principally owing to the more comfortable condition of the working classes, in consequence of the improved state of trade.							
	Barton upon Irwell	The deaths in my district have been about the average number.							
MANCHESTER	Ancoats	Total deaths 318—a general average. 111 died within the first two years of age. Of Scarlatina 3, Measles 20, Inflammation of Lungs 29, Inflammation of Bowels 8, Consumption 39, Fever 12, Croup 12, Childbirth 4, Dropsy 13, Convolsions within first month 16, ditto within 3 months 21, ditto 1 year 9. The remainder from general causes.							
• • •	. St. George	The number of Deaths this quarter 261, is rather above the average; the 10 principal causes have been—Consumption 42, Asthma 24, Pneumonia 24, Convulsions 20, Debility from Birth 14, Hydrocephalus 11, Croup 11, Hooping Cough and Pneumonia 8, Old Age 8, Coroner's Inquests 7, Other Diseases 92, Total 261; of these 78 died at 1 year and under, 33 at 60 and upwards. The average being 22 years, 3 months, and 64 days to each individual.							
9000	Market Street	354 deaths registered. Independently of the workhouse, infirmary, and coroner's inquests, the deaths in my district in the quarter ending 31st March are 150, being 19 less than the corresponding quarter of 1843.							
006%	London Road	The number of deaths registered during this quarter is less than that of the corresponding quarter for 1843 by 22, and than that of the average quarterly returns of the last year by 34. The principal decrease has been in <i>Endemic</i> , &c. diseases, excepting <i>Fever</i> , which has increased above 3 per Cent. As, however, the fever ward for the whole town and neighbourhood is in the London Road district, the difference is not great. The increase generally is in diseases of the Digestive Organs, Infantile Debility (debility from birth), and Old Age.							
****	Deansgate	Of the 242 Deaths, 104 were of Children under 5 years of age, viz.—Under 12 months 50; more than 12 months and under 2 years 29, more than 2 years and under 3 years 13; more than 3 years and under 4 years 9, more than 4 years and under 5 years 3, total 104; being very much below the average rate of infant mortality in this district.							
	Failsworth	There is an increase of deaths this quarter in consequence of Typhus, Measles, and Hooping Cough prevailing. The cases that are fatal are as follows:—Typhus 2, Measles 2, Hooping Cough 3, Coroner's Inquests 2 from Burning, Scarlatina 1.							
SALFORD	Greengate	About 35 below the average. Number of deaths in January 71, February 79, March 65, Total 215.							
	Pendleton	About the average of some quarters past.							
ASHTON	Ashton Town	The deaths this quarter are upwards of 21 per cent, less than the average of the corresponding quarters since the commencement of the registration.							
		CAUSE OF DEATH.							
		m m							
		ough lains s convulsions Birth nqueets							
		Cough halus as a a a a a a a a a a a a a a a a a							
		Dig C  Or C  Say  Say  Say  Say  Say  Say  Say  Sa							
		Hooping Cough Croup Diabetes Dysences Cholera Hydrocephalus Applexy Paralysis Convulsions Epilepsy Laryngitis Preumonifis Coolexia Consumption Philisis Philisis Philisis Philisis Consumption Preumonifis Preumonifis Preumonifis Coolexia Consumption Preumonifis Preumonifis Coolexia Coolexi							
		Male							
		Female 2 2 1 5 1 9 3 2 2 7 4 12 5 5 1 1 1 1 5 1 8							
		Surgeon 1 2 1 1 6 1 1 12 1 1 4 7 2 3 1 1 2 8 6 13 3 5 1 1 1 4 6							
		No Attendance } 1 2 1 9 2							

The state of the s	
SUP. REG. DISTRICT.	In this district the mortality is avidently despisation which may be deshrited for by
ASHTON Knott Lanes	In this district the mortality is evidently decreasing, which may be accounted for by the cheapness of provisions and the better employment of the people. The deaths registered by me this quarter are 20 less than in the corresponding quarter of last year, and 5 less than that of the preceding year. The mortality has been principally amongst children; 15 have died under 12 months, and 3 under 3 years, being the half of the whole number registered this quarter. Influenza has been prevalent but not fatal.
and the second s	The number of deaths is 29 less than in the corresponding quarter of 1843; 32 have died from Phthisis, I Influenza, 7 Pneumonia, 2 Scarlatina, 10 Asthima, 1 Nervous Fever, 1 Measles, 1 Cynanche Trachealis. Of 15 cases of Coroner's Inquests, 1 Died in Labour for want of skilful assistance, 1 Accidentally Overlain, 2 Burnt to Death, 4 Natural Deaths, 1 Accidentally killed at the Manchester and Sheffield Railway, 1 Scalded to Death, 1 Injury to the Head from an accidental fall, 1 Injury to the Spine from an accidental fall; 1 Found Drowned, 1 Accidentally Drowned; 1 Killed by a falling in of the roof of a soal mine.
	My district is free from any epidemic.
£ .	The deaths this quarter are above the average. Measles has been very prevalent.  The number of deaths this quarter is under the average; there is no epidemic prevailing at the present time. Out of a population of 8000, only 6 deaths occurred during the month of March.
Royton	Of 52 cases, 44 have been attended by qualified medical attendants, and 8 have been without medical attendance. Typhus, Meastes, Small Pox, and Varicella have all been prevalent. One fatal case of Small Pox in a family, where 4 were attacked at the same time, all without previous vaccination. A more severe and fatal attack occurred in the same locality, but in the Chadderton district.
Programme and the second secon	An unusual number of deaths from Small Pox has taken place, namely 7, all within a a short distance of each other, at the small village of Ryley Clough, with one exception, which was at Street Bridge, in the immediate vicinity, both low damp situations. No other peculiar causes of death have been more than commonly rife.
	Corresponding quarters, 1841, 92; diffto, 1842, 100; 1843, 75.  The number of deaths this quarter is 10 less than last quarter, but equals the average of any quarter in the year. I have no epidemic disease in my district this quarter.
HUDDERSFIELD Huddersfield	Sixteen above the average; 9 Coroner's Inquests out of the number.
Almondbury	Although the number of deaths in my district exceeds the last quarter by 26, yet there has been no particular disease prevalent.
. Kirkburton	There is an increase of 17 deaths this quarter above the last, and 15 above the corresponding quarter of last year, owing chiefly to young children, of which I have 22 under I year of age, 10 of Convulsions, 4 Premature Births, 4 Hooping Cough, and 4 Pneumonia.
Honley	In the corresponding quarter of 1843 there were deaths 33, in this 32. No epidemic disease has prevailed in this district. There have been 7 deaths by Consumption.
Meltham	The number of deaths registered in this district during the quarter has been rather above the average (although much below that of the corresponding quarter of last year).  Typhus has been very prevalent, and in many cases fatal, especially among the poorer classes.
Lockwood	The mortality this quarter is above an average, although there are 3 less than in the corresponding quarter of last year. 10 cases of Phthisis, principally males. Convulsions and Pneumonitis have proved fatal to several infants; there is also I case of Hooping Cough, and 1 of Croup, but I am not aware of any other epidemic prevailing at the present time in my district.
Slaithwaite	There have been fewer deaths in my district this quarter than in the corresponding quarter for the last 5 years. The only epidemic we have had this quarter is Typhus, which has been confined to a small portion of the district, and only 2 deaths from it. Of the 31 deaths, 10 have occurred in children under 10 years of age, and 9 in old people above 60 years of age.
Golcar	There is an increase of 32 deaths this quarter above the last, and 16 above the corresponding quarter last year, yet there is not any particular disease prevailing in my district.
HALIFAX Halifax	The number (192) is about 20 more than the average number of the 6 preceding corresponding quarters, and the excess is owing I believe to <i>Hooping Cough</i> .
Brighouse Brighouse	On the whole the number of deaths this quarter is not more than an average, but the mortality in children is great, 16 of the 36 being under 1 year, and 7 under a month. I attribute much of this to the neglect in parents procuring medical aid, some of them having none and others only the assistance of druggists.
Sowerby	The number of deaths registered for the quarter is above the average, but no epidemic disease has been prevailing. The number of deaths from <i>Phthisis</i> (14) and <i>Convulsions</i> (11) being the greatest of any other complaints.
	The mortality in this district, which this quarter is one-half more than last, has been principally in infancy and old age; out of 34 deaths (being the total number for the quarter), 11 have died under 2 years old, 4 above 60, 3 above 70, 2 above 80, and 1 above 90 years old.
Northowram	The number of deaths registered in my district this quarter is above an average, many having died from old age. Hooping Cough has also been very prevalent amongst children, to which the increase is mainly attributable.
Ovenden	The deaths in my district exceed those in the December quarter by 15. 11 persons have died of Consumption and 10 of Inflammation this quarter.
BRADFORD East End	An increase of 13 deaths this quarter, all from Small Pox. I have had no Typhus or Measles. The following list will shew the number of deaths from Small Pox, and also the necessity of vaccination.
	MALES.  1 Aged 38 Yearsnot previously Vaccinated—3 of this Person's Children had Small Pox, and 1 is since dead.
	25 Vaccinated 22 Years before having the Small Pox. Aged 3 Years, not previously Vaccinated Aged 3 Years, not previously Vaccinated
	5 5 7 9 9 9 8 Months. Vaccinated 9 7 7 9not previously Vaccinated
	Aged 6 Yearsnot previously Vaccinated  Aged 3 Yearsnot previously Vaccinated  18 Months  Total., Males 10, Females 6.—Not Vaccinated 14, Vaccinated 2.
	- The state of the

SUP. REG. DISTRICT.	REG. DISTRICT.	
BRADFORD		I have above 20 more deaths than usual this quarter, but it is chiefly caused by Accidents and Consumption.
	Cleckheaton	There is above an average number of deaths this quarter, Scarlatina having prevailed in some parts of my district in the months of January and February:
	Adle	This return is nearly one-third above an average; but there has not been much disease which may be called epidemic, except Malignant Sore Throat, 3 cases of which have proved fatal. I know of no cause for this increase, unless it be the sudden change of weather which took place at the end of January, from very mild to extreme cold, accompanied with keen frost and much snow, as Professor Liebig has proved, and experience corroborates the fact, that more foed as well as clothing is necessary in cold weather to keep the body in its Normal state—a condition which many of the working classes are not always prepared to comply with.
	Manningham	Though nothing like an epidemic has been observed, the number is 40 per cent. above the average.
	Thornton	Above the average, owing principally to the prevalence of <i>Phthisis</i> and <i>Pneumonia</i> .  4 Inquests and 4 deaths from <i>Typhus</i> have taken place in my district this quarter.  Rather above the average; 2 cases of <i>Typhus</i> , 1 of <i>Croup</i> , and 1 of <i>English Cholera</i> .
************	Wilsden Drighlington	The number of deaths is rather above an average, which may be accounted for by the
	Pudsey	The number of deaths in this district, during the last quarter, has been greater than during any previous quarter, with the exception of one, since the commencement of the Registration Law, being almost 40 per cent. above the average, but without any assignable cause arising from the prevalence of epidemic disease.
	Hortun	The number of deaths is greater for the last quarter than any preceding one, attributable solely to an increased Population in this district.
LEEDS	North	The number of deaths registered this quarter is 244, being 16 less than in the corresponding quarter of last year, but rather more than an average of the 4 quarters of 1843. The number of deaths, however, last year, was less by about 100 than in several previous years; and, consequently, the number registered this quarter is below the average of former years. Small Pox and diseases of the Throat and Chest have been the most fatal. Pneumonitis and Bronchitis have taken off many children and aged people, in different parts of the district, whilst Small Pox has been exceedingly prevalent in one part of it only. The locality referred to is called Newtown being situated at the Northern extremity of the Township; and from its elevation and distance from the more densely populated parts, is generally regarded as exceedingly healthy. The total number of deaths from Small Pox is 13, the greater part of them having occurred at Newtown. A very large proportion of the deaths are, as usual, those of children. Many infants die from neglect of the parents. In very many instances drugs are administered, and no regular medical aid procured till it becomes unavailing. 122 died under 5 years of age.
	F7 7 4 13	The Deaths are registered as having been caused as under:—Bronchitis 15, Pneumonitis 34, Hooping Cough and Pneumonia 19, Meastes and Meastes and Pneumonia 10, Asthma 8, Small Pox 13, Hydrocephalus 12, Debility 6, Premature Birth, 7, Cephalitis 4, Dropsy and Diseased Heart and Dropsy 7, Diseased Heart 3, Typhus 2, Croup 2, Phthisis 20, Tabes Mesenterica 12, Cancer 6, Scarlatina 2, Diarrhea 3, Apoplexy 5, Enteritis 2, Ædema 2, Old Age 6, Monomania 1, Stricture of Pyloras 1, Traumatic Tetanus 1, Convulsions 13, Erysipelas 1, Hepatitas 1, Hydro-pericardim 1, Myelitis 1, Melcena 1, Cirrhasis 1, Ovarian Dropsy 1, Hanging 1, Poisoning 1, Burning 1, Scalding 1, Rupture of the Aorta 1, Want of Food, 1. Two deaths are from Causes unknown, and the remainder from Jaundice, Peritonitis, Fracture of the Hip Joint, Diseases of the Spine, Epilepsy, Paraplexy, Hydrothorax, &c.
KINGSTON UPON	Kirkstall	Nine deaths from <i>Small Pox</i> (natural); 20 deaths above 60 years old: Males—one 67, one 69, one 74, one 75, one 77, one 80, one 82 and one 83; Females—two 62, one 63, two 66, one 72, two 74, one 75, one 77, one 95, and one 102.
HULLYORK	\{ Humber	<ul> <li>Small Pox and Measles have prevailed during this quarter in this district.</li> <li>The number of deaths is about the same as the March quarter 1843. There have been 6 deaths from Natural Small Pox, 1 six years after Vaccination, and 4 from Scarlatina.</li> </ul>
TYNEMOUTH	Tynemouth	The number of deaths registered by me during the quarter ending 31st March, 1844, viz. Males 39, Females 58—97. The average number of deaths in my district per quarter over the 6 years ending December 31st, 1843, is 79·08. The average of the 6 preceding quarters corresponding with the present is 73¹/6. The average of the 6 preceding quarters corresponding with the present is 73¹/6. The present return shews an excess of 18 above the average in the first above-mentioned case, and an excess of 235/6 above the average in the second. 39 children under 7 years of age have died this quarter, chiefly from Hooping Cough, Pneumonia, and Measles. The rate of mortality on each of the 6 years is as follows: viz.—in the year 1838 the rate per cent. was 2·1—in 1839, 2·52—in 1840, 2·01—in 1841, 2·19—in 1842, 2·27—in 1843, 2·11. The average rate per cent. of mortality over the 6 years ending as above was 2·2 on the amount of the Population.
	North Shields	Of the whole number of deaths this quarter, 44 are children under 10 years of age viz.— 6 Hooping Cough, 3 Measles, 2 Scarlatina, and 33 from General Causes, Premature Birth, General Debility, &c. The Hooping Cough and Meusles have been very prevalent, though not very fatal.
	Blyth	There have been about the usual number of deaths this quarter; there has been no epidemic disease.
NEWCASTLE UPON TYNE	Byker	One-sixth below an average of the corresponding quarters in past years, and one-eighth below the general average.
CARLISLE	St. Mary	The number of deaths registered this quarter is considerably above the average, being 49 more than the corresponding quarter of last year, which is attributable to Measles and Hooping Cough, being very prevalent and fatal, there being 49 deaths from the above causes; whereas, in the corresponding quarter of last year there were only 1 of Measles and 1 of Hooping Cough.
	St. Cuthber	This is considerably above an average quarterly number of deaths; but there does not seem to be any infectious disease prevalent to account for the increase. The deaths seem to have been principally amongst old people and young children—14 persons being upwards of 70 years each, and 22 under 1 year, which will account, in a great measure, for the increase.
between the second seco	Stanwix	The number of deaths is above the average, owing to the prevalence of Measles.  There was 1 case of Hooping Cough, also 2 of Croup. None of these have of late years been prevalent in this district.
KENDAL	Wetheral	Hooping Cough and Measles have been very prevalent in my district this quarter.  The number of deaths this quarter is more than usual, but there is no particular dis-

SUP. REG. DISTRICT. REG. DISTRICT.	
KENDAL Grayrigg	The number of deaths is within one double of last Quarter, although 3 below the average of the whole of the Quarters since the commencement of the Registration. Measles have been very prevalent in the southern part of my district, but only 1 case has proved fatal. Last Quarter the Births were to Deaths $4\frac{1}{2}$ to 1; the present Quarter they are 2 to 1.
Milnthorpe	The number of deaths is a little below the average of the corresponding Qua ters of the 6 previous years, and the district is very healthy.
ABERGAVENNY Aberystruth	The number of Deaths in my district is above the average of former Quarters, owing to the prevalence of Scarlatina.
Tredegar	The deaths are above the average occasioned by Scarlatina prevailing.
Blaenavon	Of the 30 deaths registered this Quarter, 4 are from Scarlatina and 1 from Typhus, about the beginning of the Quarter: At this time my district is healthy.
	The much increased average of Deaths is to be attributed to Small Pox, Scarlatina, and Coroners' Inquests.
Lower	43 cases of Scarlatina, 36 of Phthisis, 3 of Small Pox, of which one had been Vaccinated, 4 of Puerperal Fever, and 28 of Fits, though from what causes it is impossible to say.
Gellygare	In this Quarter there are 52 deaths—a greater number than the average from the prevalence of the Scarlatina in my district.
NEWTOWN AND   Newtown	The deaths this Quarter have been rather more than the preceding Quarter, chiefly from Old Age and Decline; a few deaths by Measles.
	There is no epidemic prevalent in my district; most of the number died from Old Age and Decline.
WREXHAM Hope	The number is above the average ; $Typhus$ prevailed in my district—4 fatal cases in one house,
Malpas	The number of deaths in this Quarter is 44, being 3 more than in the preceding Quarter and 3 less than the corresponding Quarter of 1843. No epidemic prevalent during the Quarter.
Wrexham	The number of deaths registered this Quarter is 82, being 21 less than the corresponding Quarter of 1843, and 9, 9, and 15 respectively above the 3 preceding Quarters;—8 have died from Phthisis, 4 Tubercular Phthisis, 11 Convulsions, 11 Old Age, 5 Bronchitis, 3 Asthma, 2 Influenza, 2 Scarlatina, 2 Typhus 3 Pneumonitis, 2 Hydrocephalus, 1 Tabes Mesenterica, 1 Chronic Diarrhoa, 2 Paralysis, 2 Fever, 1 Jaundice, 1 Chronic Bronchitis, 1 Disease of the Heart, 1 Puerperal Fever, 1 Trismus, 4 Coroner's Inquest cases, 2 Premature Birth, 1 Disease of Womb, 1 Effusion of Serum at the base of the Brain, 1 Debility after Abortion.
HOLYWELL Holywell	The prevailing complaints have been the 3 following, viz.—Consumption 18, Convulsions in Children 15, Scarlatina 14, Deaths of aged Persons from 70 to 96 years, 11, Inquests 7, other causes 34—Total 99.
	***************************************
GENERAL REGISTER OFFICE, 2 May, 1844.	
2 2009, 2021	

### A QUARTERLY TABLE

OF THE

# Mortality

IN A

## HUNDRED AND FIFTEEN OF THE DISTRICTS

OF

# ENGLAND

(INCLUDING THE PRINCIPAL TOWNS,)

SHEWING

THE NUMBER OF DEATHS REGISTERED IN THE SIX YEARS 1838—43, THE AVERAGE

NUMBER OF DEATHS IN THE FIVE SPRINGS 1838—1842;


AND THE NUMBER OF DEATHS IN THE SPRING QUARTER 1844, ENDING JUNE 30TH.

PUBLISHED BY AUTHORITY OF THE REGISTRAR GENERAL.



LONDON:

HARTNELL, PRINTER, RED LION COURT,



### TABLE

OF THE

### MORTALITY

In 115 of the Districts of England (including the principal Towns): shewing the Number of Deaths Registered in the 6 years, the average Number of Deaths in the 5 Springs, 1838—42, and the Number of Deaths in the Spring Quarter of 1844, ending June 30th.

	Popu-	De	aths R	Pariete	red in	the ve	ars	1838-	-42	Spring ding 1844		Ponn	n	eaths I	Register	ed in t	he vears		1838-	-42	Spring ding 1844
DISTRICTS	lation	100	ntio y	egiate	104 111	me je		Quar * Ave		in the ter, end 30th,	DISTRICTS	Popu- lation							Quart • Ave	rage	in the ter, end 30th,
	1841	1838	1839	1840	1841	1842	1843	Of Five Years	Of Five Springs	Deaths i		1841	1838	1839	01810	1811	1842	1843	Of Five Years	Of Five Springs	Deaths i
•Metropolis					1						N. Midland division										
West Districts	301326	8114	6538	6936	6599	6826	7212	1751	1738	1802	Leicester	50932	1170	1291	1730	1358	1457	1255	350	326	341
North Districts	366303		3	8594	8261	8550	9038	2112	2004	2191	Lincoln	36110	620	731	817	, 800	720	786	184	194	180
Central Districts						. 1		2395	2255	2044	Nottingham	53080	1209	1540	1794	1239	1429	1393	361 310	315 322	327 359
East Districts	393247 479469					1		2640	2417	2433	Basford	59634 35015	769	1200	1549 926	914	1245 909	1216	217	208	
	1915104					··· · ,		$\frac{3044}{11943}$	2686	$\frac{3001}{11471}$	N.Western	93019	705	002	02.7	713	505	322			
S. Eastern				-				11343	11101	11411	division Stockport	85672	2410	2591	2574	2048	2073	2203	585	<b>63</b> 0	418
division  Maidstone	33210	737	726	700	648	815	774	3.03	104	202	Macclesfield	56018	1315	1528	1745	1419	1403	1551	370	394	338
Brighton	46742			980	1102		1117	181 251	$\frac{184}{230}$	219	Gt.Broughton,	49085	1040	1237	1384	1246	1189	1091	305	315	237
Isle of Wight	42547			737	891	721	703	182	193	191	inc. Chester 5	223054	6627	7435	8470	7556	7407	7458	1875	1744	Ę.
Portsea Island	53036			1140	1211	1152	1342	297	271	363	West Derby, }				2206	2202	2405	2312	509	455	
Winchester	23044			526 402	416 395	469 394	471 314	116	113 94	117 98	(adj. Liverpool)	88652	1625	1746							
Windsor		381	367	402	990	994	314	97	J-±	30	Blackburn Preston	75091	1612	2190	2140	1692 2026	1665 2050	2031 1938	465 513	492 502	1
S. Midland division											Rochdale	60577	1726 1477	1812	1868	1417	1476	1531	396	426	
St. Albans	17051	378	310	331	300	340	291	83	89	73	Bury	77496		1990	2170	2064	1987	1832	506	543	E
Wycombe	34150	788	692		751	679	748	187	210	<b>22</b> 9	Bolton	97519	2432	2857	2900	2572	2607	2596	668	672	7
Oxford	19701		1 .	406				102	94	2	Wigan	66032	1	1912	2144	1561	1588	1832		447	
Northampton	28103 31767			670 743		622	677 762	168 171	165 172		Prescott	43739		1056	1155 2399	997	1075	920 2529	260 591	272 535	
Cambridge	24453			639				144	144		Manchester	93736		2418 6774	6489	2431 5821	2427 6151	6263		1543	
											Salford	70228		2449	1998	1971	1977	2098	FOO	519	2 4
Lastern division											Ashton	173964	į	5489	4873	4282	İ	4391	1196	1377	993
Colchester		1	}					113		5	Work										
Ipswich	25254 61846	1	\$					150	100		division							000	F00		101
Norwich	24031			1752 504	493	1000		396 118		0 1	Sheffield	85076	Į	,	2410 2243	2231	2182			1	8
Yarmouth		24,	310	001		910	020			1	Huddersfield .	107140		2080 2402		2122	1	1			
S. Western division			-								Bradford	132164	1		3176	3041	1			3	
Devizes	22130	200	510	481	429			LLO	111	118	Leeds	168667	4289	4388	4488	4360	4615	4329	1		3
Dorchester	23380		004			1	524	10,	1	1	\$	41130	1060	1023	1442	1206	1	i	0.77		
Exeter	31333 47103	1		1	0.00			1 202	1	3 10 0	1	47779	876	1217	1209	1039	1133	1048	274	292	267
St. Thomas Plymouth	36521	1	-		800		1		1		Northern									-	
Redruth	4806	1	1	1	00-	4		1			CLA LEGICAL	56226	1479	1524	1513	1512	1357	1247	369	37-	307
Penzance	50100	762	848	976	1	1	996		1		Gateshead	38747			1016		1	§	000		
Bath	6923	2 1567	1602	1974	1728	1598	1624	423	436	433	Tynemouth	55625	1112	1285	1243	1340	1327	1229	315	322	268
Western division				-							Newcastle-on-	71850	1829	2117	1957	2104	1726	1922	487	469	373
Bristol	64298	1845	1675	2330	1895	1771	1742	476	462	424	Carlisle	36084	855	739	901	925	840	762	213	193	199
Clifton	66233		2230	2				1	1	427		35676			696	674	597	}	1 200		
Stroud	38920	1	000						1		2	34694	729	825	821	601	714	666	184	198	188
Chelteuham	4922 33646		1 6740		1		5	1	1		33										
Shrewsbury	21529		00,	1				1	,		I AT	50834	1150	1206	1374	1458	1109	1116	315	357	332
Worcester			013		i .		į.		L		1 m	25037	1	.615	563	648		Į.	20 400	170	99
Kidderminster	29408				1	3	1	2 40			Merthyr Tydfil	52864	t .	1246	1566	1423	1	1			
Dudley		1		1 .	1	Į.			1		Newtown	25958	ì	468	619	539	420		129	144	
Walsall		1		1	ī.					_	Wrexham	39542 40787		823 791	966	900			217	227 222	221 209
Wolverhampton Wolstanton'		-	1	4	1	è	i	6000			Holywell Anglesey	38105			864 632	800 597	738 624	815 575	7		_
Birmingham		1		2	1		ł	000		_					-						
Aston			10.58			1093		0-0	3		of the Metropolis	4663308	109321	116505	124664	114664	115834	113869	29019	29107	27454
Coventry	3102	8 68.	1 724	- 616	905	890	838	191	181	199	GRAND TOTAL	6578912	162867	162605	171694	160733	161948	163201	40992	10208	38925
																-					
\$ 100 TO T		2 .7														-	THE RESERVE				

<sup>\*</sup> The Deaths in the Metropolis for the Years 1840 1-2, have been derived from the Weekly Tables, the Returns for each Year comprising 52 Weeks, or 364 Days. The last quarter in the Metropolis ended June 29, 1844. The Returns from other Places are for the years ending December 31, and the quarter ending June 30th.

<sup>\*</sup> Wandsworth District is included in the return for the Metropolis.

### A TABLE OF THE MORTALITY

IN THE

### Metropolis,

### SHEWING THE NUMBER OF DEATHS FROM ALL CAUSES,

Registered in the 13 Weeks ending June 29th, 1844.

CAUSES OF DEATH	Quarterly Average of Deaths in the 5 Springs of 1839-43	Deaths in the Spring Quarter ending June 29th 1844	CAUSES OF DEATH	Quarterly Average of Deaths in the 5 Springs of 1839-43	Deaths in the Spring Quarter ending June 29th 1844
ALL CAUSES	10960	11471	IVLaryngitis	. 7	17
SPECIFIED CAUSES	10912	11432	Quinsey	18	23
I Zymotic (or Epidemic, Ende-)	3000	0.451	Bronchitis	130	194
mic, and Contagious) Diseases	1993	2451	Pleurisy	23	19
SPORADIC DISEASES:-			Pneumonia	813	715
II Dropsy, Cancer, and other Dis-			Hydrothorax	54	67
eases of uncertain or variable	1300	1209	Asthma	198	161
Seat			Phthisis or Consumption	1919	1838
III.—Diseases of the Brain, Spinal	1887	2006	Disease of Lungs, &c	178	195
Marrow, Nerves and Senses (			V.—Pericarditis	11	29
IV.—Diseases of the Lungs and of the other Organs of Respiration	3340	3229	Aneurism	9	14
V.—Diseases of the Heart and Blood			Disease of Heart, &c	240	335
Vessels	260	378	VI.—Teething	205	165
VI.—Diseases of the Stomach, Liver, ?	hee	0.47	Gastritis	} 193	11
and other Organs of Digestion §	755	847	Enteritis	5 100	169
VII.—Diseases of the Kidneys, &c	68	94	Peritonitis	17	29
VIII.—Childbirth, Diseases of the	108	99	Tabes Mesenterica	61	125
Uterus, &c	100		Worms	3	6
IX.—Rheumatism, Diseases of the Bones, Joints, &c	77	82	Ascites	9	17
X.—Diseases of the Skin, Cellular			Ulceration (of Intestines, &c.)	15 22	9
Tissue, &c	14	12	Hernia	22	23 32
XI.—Old Age	795	673	Intussusception	5	7
XIIViolence, Privation, and In-			Stricture	7	3
temperance	314	352	Hæmatemesis	6	10
I.—Small Pox	144	425	Disease of Stomach, &c	50	74
Measles	355 325	208 601	Disease of Pancreas	• •	1
Hooping Cough	438	361	Hepatitis	13	22
Croup	99	126	Jaundice	26	39
Thrush	49 59	42 83	Disease of Liver, &c	.99	105
Dysentery	12	18	Disease of Spleen	1	**
Cholera Influenza	5	9 <b>2</b> 8	VII.—Nephritis	5	9
Ague	25 3	-3	Ischuria	2	
Remittent Fever	6	4	Diabetes	4	3
Typhus	400	455 68	Cystitis	3	2
Erysipelas	- 8	. 19	Stone	5 6	16
Hydrophobia	.1	1	Disease of Kidneys, &c	44	59
II.—Inflammation	81	7			1
Hæmorrhage	40 411	42 333	VIII.—Childbirth	78 2	67
Abscess	42	20	Ovarium Dropsy	3	6
Mortification	55	46	Disease of Uterus, &c.	24	23
Purpura	3 29	43	IX.—Arthritis.	1	1
Cancer	106	170	Rheumatism	34	27
Tumor	19	7 19	Disease of Joints, &c	43	54
Atrophy	13 76	137		1	
Debility	239	236	X.—Carbuncle	1	2
Malformations	12 173	$\begin{array}{c} 22 \\ 124 \end{array}$	Ulcer	4	7
III.—Cephalitis		153	Fistula	5	1
Hydrocephalus	461	497	Disease of Skin, &c	4	2
Apoplexy,	209	294	XI.—Old Age	795	673
Paralysis		213 614		7	
Tetanus	4	7	XII.—Intemperance Privation	3	13
Chorea Epilepsy		2 55	Violent Deaths	304	333
Insanity		18			
Delirium Tremens	22	29	Causes not specified	48	39
Disease of Brain, &c	106	124			

# QUARTERLY METEOROLOGICAL TABLE,

Compiled from the Weekly Tables furnished to the Registrar General by the Astronomer Royal.

Deaths registered in the Metropolis from Small Pox, Measles, Scarlatina, Hooping Cough, Typhus, Diarrhaea, Dysentery and Cholera, Influenza, Consumption, and other diseases of the Lungs; the numbers at each age,\* and the total deaths't (except violent and sudden) in each of the 13 weeks ending June 29, 1844.

ı						2	- 1-1-1-1			ā					1
	610.68	10 ·	29.778	2 9.941	29,840	29,890	29,911	30.039	29.785	30,148	30,016	30,090	30.007	inches 29,807	Mean height of the 72 observations, duced to 32 deg. I
	89 89 84	83.3	79.6	83.1	74.1	69.8	66,8	74.6	74.1	67.3	74.6	8.00	66.7	67.0	Highest during the
	34.0	48.0	8.0F	17.5	43.6	42.1	43.8	34.6	44.3	35,9	88 28	96.3	34.0	37.0	Lowest during the
	65.4	69.0	70.3	74.5	70.6	\$30° \$3	63.9	63.3	67.8	61.0	0.08	61.5	1:19	64.9	Of the Highest on e
	\$0.5°	\$1.8	52.1	51.7	\$0.3	44.2	44.9	42.0	46.0	41.8	42.5	43.2	40.7	10.4	Of the Lowest on ea from 6 observation
Ī	19.0	17.2	18.2	80 80 80	20.3	14.0	0.61	\$2 •3	%1.8	19.7	23.5	18.3	20.4	22,9	Difference
I	55.0	60.5	60.8	62.7	59.5	50.9	\$2.9	82.1	56.7	51.0	54.3	05 22.	50.3	50.9	Mean of 72 observa
ľ	47,2	53.6	59.4	50.5	50,6	44.8	46.7	65.22	49.0	43.4	45.9	46.6	42.2	43.2	Mean of 72 results
	111.7	111.7	108.5	107.1	99.0	91.0	99.3	108.2	103.7	93.8	102.8	87.3	95.4	86.7	During the week
	89,6	92.4	94.6	100.0	93.2	75.8	88.6	92.7	94.1	85.7	90.9	79.0	82.4	82.7	Mean of 7 obser- vations
ľ	21.6	39.0	34.8	36.0	36.9	32.0	31.2	26.5	31.7	24.0	24.5	24.4	21.6	23.5	During the week
	37.0	46.8	\$9. 12	4 <u>1</u> 8	42.7	38.0	38.0	32.6	37.1	31.3	32.6	35.5	12 12	29.9	Mean of 7 observations
	7.8	6,9	8.4	12.2	8.9	6.1	6.2	0.0	7.8	7.5	00 4	5.7	6.1	7.7	Mean of 72 differen
	15,4	13,9	15,2	21.1	17,6	12,2	13.2	14,0	16.7	15,1	17.1	12.3	16.0	16.4	Mean of the great on each day, 6 oh vations
	5	\$100 0 0 0 0 0	60 60	ယ မိ	2,6	0.9	1.9	8	0.9	0.7	gui a a	0.6	<b>89</b>	1,3	Mean of the least each day, 6 obse tions
	+	1.1	+0.4	+3.0	+2.0	4:1	11%	O;-ii	+4.1	10.6	+6.1	+6.2	+4.1	+6.2	Difference between rature of the we temperature of t an average of 25 y
		N.N.E.	s.w.	W.S.W.	S.W.	N.N.E.	N.E.	N.	N.N.W.	E,S.E.	and and N.N.E.	S.W.	W.S.W.	S.W. and E.N.E.	General direction
	10.0	9.0	<u>د</u>	5.0	9.0	7.0	10.0	Cit En	ŧ,	3,0	3.0	3.0	35	3.0	Greatest pressure the week
	0.86	0.04	0.2	0,6	0,2	0.8	1.4	0.6	0.1	0.1	0.04	0.1	0.8	0.2	Mean for the week
	94.8	625	855	1170	1265	1220	1400	1065	740	960	690	760	900	miles 670	The amount of Ho of the air in each
	55.00	80	6,3	ab. Sea	6.5	00	6,9	5.7	6.0	3,1	3.7	6.7	3.9	2,9	Mean amount of Clo
	2.31	1.55	0.10	0.02	0.03	0,19	0.01	z.	0.06	0.02	0.05	0.02	0.26	3	Rain in inches (7 da
	425	51	20	43	49	31	36	30	27	22	2	20	21	21	Small Pox
	208	19	21	13	22	-	23	16	9	40	10	\$	222	19	Measles
	60	60	78	62	56	63	49	=	35	24	35	32	<u>د</u>	ಚ್ರ	Scarlatina
	361	13	10 10	20	16	20	27	88 80	26	39	27	40	27	46	Hooping Cough
	455	39	36	27	32	33	32	30	es es	36	38	=	39	<b>63</b>	Typhus
	110	15	12	10	7	8	60	6	13	9	బ	9	C7	Cri	Diarrhota, Dysenter
	28	0	<b>b</b> et	-	jus .	c	4	**	ω	-	co .	40	6	4	Influenza
	1838	146	125	136	152	142	136	138	134	168	162	142	136	121	Phthisis or Consum
	1391	18	108	E	97	104	103	105	107	105	93	120	121	136	Other diseases of the
	5172	445	434	397	417	395	497	397	336	373	365	363	396	428	0 to 15
	3712	271	282	286	302	284	262	273	299	318	304	270	286	275	15 to 60
	2113	158	4	149	164	172	168	166	152	154	167	176	167	179	60 and upwards
	31 10 014	875	858	802	8.84	853	857	837	788	845	836	810	850	869	Deaths from all can violent and sudden

• The Ages of 17, on account of the corrections for returns not received,
were not specified in the Table.

Inquests are held on violent and sudden deaths which are returned irregularly.

Mean, Highest, or Lowest, of the 13 weeks.

29th 22nd

23rd 1st qr. 16th, 15(1)

15th, Greatest declination N.

DOW

June

18t

May 31st full and in Perigree

7th, 3rd qr.

25th 1881

25th, 1st qr.

May

48h

and, full and in Perigree

27 tb 20th 1344

24th, 17th,

, ist qr.

Tith

9th, 3rd qr.

April

604

9th, 3rd qr. 3rd, fall Week

week

each day, ach day,

Highest in the sun

Lowest on the grass

eatest beer-it on erva.

Mean

Self-Registering

Difference be-tween the dew point temperature and air temperature

en the mean tempe-eek, and the mean the same week on years

Pressure in lbs. on the square foot WIND

in

oud, 0-10

y, and Cholera

e lungs

Deaths at Three Ages, (exclusive of violent and sudden Deaths)

uses, (exclusive of in deaths)

THERMOMETERS

Moon of the Phases

### SUMMARY OF METEOROLOGICAL OBSERVATIONS, Made at Helston, in Lat. 50° 7′ N., Long. 5° 18′ W., for April, May, and June, 1844. By M. P. MOYLE, Esq.

	By M. P. MOYLE, Esq.														
				]	BAR	ОМ	ETEF	<b>.</b>							
MONTH	9 a,m		np. of fer.	3 p.m.		np. of Ier.	9 p.m		Temp		Mean Mear		Mean of i. Temp.	Corrected to 32 deg. Fahrenheit	
April	30.21	3 5	8.1	30.215	6:	2.13	30.22	26	62.	.93	30.21	8	61.05	30.144	
May	30.20	1 6	1.5	30.191	6.	3.00	30.2	11	66	.10	30.20	01	63.53	30.116	
June	30.05	7 6	4.9	<b>3</b> 0.066	6	6.70	30.08	30	67.	.20	30.06	37	66.26	29.982	
			,		GREA	TEST	RANGE								
	Max	D	ay	Min.		Day	Rang	e	Grea rang 24 h	e in	Day				
April	30.6	0	9th	29.70		4th	<b>0.</b> 90	, .	0.	26	5th to	6th			
May	30.4	7 2	end	30.00		6th	0.47	1	0.25		16th—	17th			
June	30.3	4 1	6th	29.70	2	4th	0.60		0.	.36	17th—1	18th	;* 		
	THERMOMETERS.  COMMON IN SHADE REGISTERING														
	Greatest														
	9 a.m	. 3 p	o.m.	9 p.m.	M	ean .	Max.	М	in.	Mea	n Ra	nge	Greatest range in 24 hours	Day	
April	50.66	50.66 58.76			52	.20	59.83 4		3.70 51.		74 16.13		0.24	9th	
May	55.20		.90	51.30	56	.13	62.70 47		.00	54.5	66 -15	.70	0.26	7th	
June	60.70	67	.40	55.90	61	.33	68.60	52	.40	60.5	50   16	.20	0.24	10th	
		. 46 1 1	- 1114	CE OF V				AV	ERAG	E CLOU					
April.			.30	9 p.m. 2.70	1	ean .56	9 a.n			3 p.m. 2.3	1	9 p.m.		Mean 2.56	
May	1.20	1	.60	0.71	1	.17	2.6		,	2.5		1.8	1	2.30	
June	1.10	1 ., 1	.53	0.73	1	.14	. 3.8			3.5		3.14		3.90	
						ONS	OF V	VIN	D.					,	
_	APRII			MAY			JUNE			• 4		Wet Days	Days Observ.	Rain	
E	a.m. 3 p.m 9 7	9 p.m.	9 a.m.	3 p.m. 9	p.m.	9 a.m.	3 p.m. 9	9.m.	13	gr or majorities a	21	1 1		1 1	
S.E.	0 0	0	0	0	0	0	1 :		·A]	pril		3	30	0.22	
S. S.W.	$\begin{bmatrix} 2 \\ 6 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \end{bmatrix}$	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	0 1	0	0	17	16	16	M	а <b>у</b>		3 .	31	0.21	
W.	3 4	6	1	3	2	.1	4	4	Jı	ine		., 6	30	0.86	
N.W.	7 11	12	6	9	0	3	6	<b>1</b> 5		A 10 - 10 - 10	to be force of the same of			l some it	
N.E.	2 0	0	6	10	11,	., 3	.1 .	1,	-5	, .	-	12	91	1.29	
	FALMO	UTH	.—(D	educed	from	Obse	ervatio	ns l	y L	OVEL	L SQ	UIRE,	Esq.)		
		В	AROI	METEI	R.	4. 1			ERM	ом	ETER		Total		
1844	1 1 1 1 1 1 1 1 1	Attd.		erage		A ++A	1			verage	:	,	Rain	1844	
April	9 a.m. 30.18	Ther.	3 p.m. 30.18	I Her.	9 p.m.	I HCI.	9 a.m.		]	9 jp.m.	Máx.	Min.	inches	(A- 3)	
May		60	30.15	61.7	30.17 30.15	61				48.81	56:7	45.98		(April)	
June		, ,	30.00	-	30.00		56		1	51.53	1	48.33		(May)	
o diffe. , ,	00.00	00.0	00.00	03.2	90.00	63.8	6.3	63	.0.	56.46	65.8	04.45	2.729	(June)	

### REMARKS.

The Quarterly Returns are obtained from 115 Districts, sub-divided into 576 Sub-Districts. Thirty-four Districts are placed under the Metropolis, and the remaining 81 Districts comprise, with some agricultural Districts, the principal towns and cities of England. The population was 6,578,912 in 1841.

The Deaths registered in the last quarter (ending June 30th) amounted to 38,925, which is less by 7116 than the Deaths in the previous quarter, and 1283 less than the average of the corresponding Spring quarter in the 5 years (1838-42). Allowing for the increase of the town population, the mortality was 10 per cent. below the average of the season.

The reduction in the mortality has been unequally distributed over the kingdom; but as a general rule, it has been most remarkable in the large manufacturing Districts.

	Average of 5 previous Spring Quarters, 1838-42	Deaths in the Quarter ending June 30, 1844.
Manchester	1543	1260
Salford	. , 519	417
Ashton and Oldham		993
Leeds	1107	936
Sheffield	. 577	464
Stockport	630	418
Liverpool	1744	1547

Birmingham remained the same; and in a few of the other manufacturing Districts, to be shortly mentioned, the mortality was above their average.

Small-pox and Scarlatina have been epidemic in the metropolis; and towards the end of the quarter, 51 persons (chiefly children) died of the former, 78 of the latter disease in one week. The Deaths in the quarter from Small-pox were 425. 107 of the number were reported by the informants to have been vaccinated; but the evidence of this would necessarily be, in many cases, uncertain and imperfect. Influenza has declined; and since the mean temperature has risen above 60 degrees, bowel complaints have become more prevalent.

The epidemics most frequently mentioned in the country Districts are Scarlatina, Measles, Small Pox, Hooping Cough, and Typhus.

The Quarterly Meteorological Table has been compiled from the Weekly Tables supplied by the Astronomer Royal. The mean temperature (55° 0) was 1° 9 above the average of the season; the highest temperature of the air (observed) was 83° 3, the lowest 34° 0; the thermometer once rose in the sun to 111° 7, and fell on the grass so low as 21° 6. The mean dew point was 47° 2. The quarter has been remarkable for the small quantity of rain which fell. The rain collected in 13 weeks was only 2.31 inches; and of this, 1.55 inches fell upon the last week of June.

The results of some Meteorological Observations by M. P. Moyle and Lovell Squire, Esqs., are given in another page. The publication of the results of a series of simultaneous Meteorological Observations, made on an uniform plan in different parts of the kingdom, could not fail to be highly interesting.

Districts in which the Mortality was HIGHER than the average Mortality of the Spring Quarter in the same Districts.—The South Districts of the Metropolis, Portsea Island, Northampton, Bedford, Clifton, Stroud, Hereford, Kidderminster, West Derby, Wigan, Bradford, Merthyr-Tydfil.

Districts in which the Mortality was lower than the average Mortality of the Spring quarter in the same Districts.—Brighton, Oxford, Norwich, Dorchester, Bristol, Worcester, Wolverhampton, Stockport, Macclesfield, Great Boughton (Chester), Liverpool, Preston, Bury, Bolton, Prescott, Manchester, Salford, Ashton, Sheffield, Huddersfield, Leeds, York, Sunderland, Tynemouth, Newcastle-on-Tyne, Pontypool, Newtown, Anglesey.

## REGISTRARS' NOTES TO THE JUNE QUARTERLY RETURN.

The following Direction was addressed to the 447 Registrars in the Country:-

"If at any time the number of deaths registered during the Quarter has been above the average, state, in a note at the foot of the Return, whether any epidemic disease, such as Measles, Typhus, &c., has been prevailing in the District; or if there be any other known circumstance which will account for the increase."

In compliance with the above direction, 181 Registrars have appended explanatory notes to their Returns; the remaining 266 Registrars apparently observed a decrease, or no obvious increase, in the mortality.

An Abstract of the Notes to the Returns is subjoined.

	DEC DICEDICE	
sup. reg. district.  MAIDSTONE	REG. DISTRICT.	The Deaths during this quarter have been considerably above the average; Scarlatina
MAIDSTONE	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	has been very prevalent amongst children under 14 years of age, 20 Deaths from that disease having been registered.
	Loose	The number of Deaths is rather above the average this quarter; Scarlatina has been unusually fatal among children in one part of my district.
BRIGHTON	The Palace	Only once since the Registration Act passed has the number of Deaths been so few in this district as during this quarter.
	Kemp Town	The return is 17 less than the corresponding quarter, 1843, and 36 less than the previous quarter; the district continues in a very healthy state.
ISLE OF WIGHT	Newport	Quarterly average number of Deaths in this district for the 7 years ended 30th June, 1844, 67; quarter ended June 30th, 1844, above the average 11; Measles prevalent during the quarter, 12 cases fatal.
PORTSEA ISLAND		Of the Deaths registered this quarter 13 were from Measles and 9 from Small Pox; 7 of these without previous vaccination and 2 after vaccination.
<u> </u>	Kingston & Landport	The Deaths registered this quarter exceed those of the corresponding quarter of 1843 by 4; in the quarter ending June, 1843, I registered Measles, I; Small Pox without vaccination, 1; Pneumonia, 3; and during the past quarter Measles, 5; Small Pox without vaccination, 7; after vaccination, 2; Pneumonia (chiefly after Measles) 8; being an increase in the epidemical diseases among children of 17, over that of 1843.
WINDSOR ,	Egham	Rather above the average of corresponding quarters, district generally healthy; there has been one case of Small Pox (natural.)
ST. ALBANS	St. Albans	The number of Deaths during the past quarter is below the average.
WYCOMBE	High Wycombe	This quarter's return is above the average, being 17 more Deaths than in the corresponding quarter of 1845, owing to <i>Measles</i> followed by Inflammation, which has been severe and proved fatal in many cases in my district.
	West Wycombe	The number of Deaths this quarter is above the average, Measles prevailing.
	Great Marlow	The number of Deaths is one less than the last quarter, Influenza has been very prevalent 4 cases having proved fatal. The number of Deaths of old people has been very large this quarter, they are as follows:—Il above 60, 9 above 70, 4 above 80, 1 above 90, with the exception of Influenza my district is very healthy.
NORTHAMPTON	All Saints	This is the largest number of deaths in my district in any one quarter during the whole period of the Registration. Including the present return of 100 deaths, the average of the same quarter for the 7 years is about 67½, consequently there is an increase of nearly 50 per cent. The large number of children who have died of <i>Measles</i> will account for the difference.
	St. Giles	The mortality this quarter is less by 8 than the preceding one, I have had 22 cases of Measles and 3 of Small Pox this quarter.
BEDFORD	,	Number of deaths rather above the average, principally owing to Measles being very prevalent in this district during the quarter.
	. Bedford & Kempston 👡	The number of deaths registered this quarter is about one-sixth less than the average number registered in the corresponding quarters of the preceding 6 years. Measles and Hooping Cough have prevailed as epidemics over a considerable part of this district, but the mortality has been unusually slight; this is most probably to be attributed to the fine dry weather. Measles commonly occurs as an epidemic in the town of Bedford every seventh year, mostly in the winter, and it has been in general very fatal. It has, however, on this occasion anticipated its ordinary period of recurrence by about six months.
CAMBRIDGE	. St. Andrew the Less	The number of deaths registered in this district during the quarter is 62, which is rather above the average (although much below that of the two last quarters) 8 children have died from Small Pox without previous vaccination, Many parents have a prejudice against vaccination here, and Small Pox lingers in this district.
	. St. Giles	There are four more deaths registered this quarter than in the corresponding one of last year. There are eleven deaths from Small Pox, only one of which was after vaccination.
IPSWICH		Out of the 49 deaths registered, 13 died of Consumption, 3 of Hooping Cough, 2 of Scarlatina, and 3 of Pneumonia.
CDD 45 Transcription	St. Matthew	Rather above the average owing to Hooping Cough having been prevalent and fatal.
	. Ivorthern	The number of deaths is 9 above the corresponding quarter of 1843. There has not been any particular disease prevailing in my district.
DEVIZES	. Lavington	The number of deaths this quarter is under the average, district generally healthy.

SUP. REG. DISTRICT	REG. DISTRICT.	
DEVIZES	Bishops Cannings	The number of deaths this quarter is 7 more than the corresponding quarter of last year, the district has been very unhealthy during the quarter, Fever and Consumption have prevailed in the district.
		Rather below the average, district very healthy.  Below the average—of the 72 registered, 14 were above 70 years of age, and 20 died of
EXETER	St. David	Consumption
tr <del>anscolut, il como</del> gipti pre la prese più p	St. Sidwell	Exceeding the corresponding quarter of 1848, by 29; 4 cases of Puerperal Fever, 40 of Consumption and Inflammation of the Lungs and Air Tubes, 2 cases of natural Small Pox without vaccination occurred about 6 weeks since, but the disease does not appear to have spread.
S7. THOMAS	Exmouth	An increase of deaths as compared with the last quarter, and the corresponding quarters of last year, 5 deaths from Phthisis, and 5 from Hooping Cough.
	St. Thomas	36 being less than the number registered in the corresponding quarter of either of the four preceding years; the present return includes 1 case of Typhus, 1 of Rheumatic Fever, 2 of Hooping Cough, 2 Erysipelas, and 2 Inquests.
suspenses and the second secon	Kenton	Hooping Cough and consequent Inflammation of the Lungs, have been very prevalent in April and May. The number of deaths last quarter were 22, including 3 cases of Sudden Death.
diversited and action families dispressing a series of	East Budleigh	The number of deaths this quarter is unusually small compared with the amount of sickness that has prevailed.
- International Control of the State of the	Heavitree	The return this quarter presents nothing extraordinary; deaths from disease of the Lungs and Heart are in greater numbers than last quarter, which I attribute to the weather—wind from the east with hot sun.
- minimum and anti-all-material and a second a second and	Woodbury	This return is above the average, but no particular cause can be assigned for it; 3 males and 1 female of the return died at the age of 76, 79, 89, and 92.
PLYMOUTH	Saint Andrew	The mortality this quarter is rather below the average of the Spring quarters, although Small Pox has been prevalent and also the Measles.
e and a street a		Small Pox has prevailed considerably; 10 deaths from that disorder have been registered, the whole of which were unvaccinated; 11 cases of Phthisis, 4 of Pneumonia, and 4 of Bronchitis occur in this quarter, there have also been 4 violent deaths registered.
REDRUTH	Camborne	The number (49) is about the average.
PENZANCE	Saint Just	The number of deaths this quarter being 58, is 15 above the average, reckoning from the commencement of the Registration Act; the cause of which excess appears to have been the prevalence of the Hooping Cough among children; as, out of the 53 stated, 28 have occurred from that disorder. There has also been one case of the Measles.
engrippercontengengengengengeng A.O. O. D. A. O. S.A.O. O.	St. Buryan	There is an increase of 3 on the corresponding quarter of last year; no particular malady has been fatal. Hooping Cough and Measles are very prevalent but not very fatal.
Summarine Suspendente transformer p & 6 C. C. C. C. C. C. C.	Marazion es en este est est est	There have been less deaths in the past than the preceding quarters; Scarlatina has disappeared, but Measles prevails to a great extent, but no cases have as yet proved fatal.
BATH	Lansdown	87, this is above the average of the quarter; Scarlatina caused 9 deaths, Hooping Cough 7, Measles 2.
\$ 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,	Walcot	My district is particularly healthy; the smallest amount of deaths in this district since the commencement of the Registration Act.
	Batheaston	11 deaths from Consumption, 4 children from Scarlatina, and 3 from Hooping Cough, 8 Coroner's Inquests.
s era evererarandinara	Twerton	This quarter I have registered 15 or 16 above the average, 7 of the cases were Scarlatina 2 Typhus and 6 Inquests.
BRISTOL	Castle Precincts	Of the number 3 died of Small Pox without previous vaccination, 2 of Hooping Cough 2 of Scarlatina, and 1 of Typhus.
To the second se	St. Paul	The number of deaths has been above the average; Small Pox has prevailed during this quarter, 10 fatal cases have been registered in this district; 8 without previous vaccination, and 2 in which vaccination was said to have been performed.
CLIFTON	Clifton	Rather above the average, <i>Hooping Cough</i> has been very prevalent, 20 deaths have been registered this quarter from the above disease, and 1 from <i>Small Pox</i> without previous vaccination.
	St. Philip and Jacob	Twenty-six of the 142 deaths, caused by Small Pox, which is extremely prevalent in this district.
9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	Ashley	The number of deaths registered during this quarter is greater than any yet registered by me except the previous quarter, which only exceeded it by one. It is nearly double the average number of the corresponding quarter. Epidemics have been registered from Scarlatina, 3 from Hooping Cough, 2 from Measles, and 1 Small Pox.
constitutionalist dissentation g g g g g g g g g g g g g g g g	St. George	The number of deaths registered this quarter is increased by 10 Inquests held in preceding quarters, this will leave the mortality this quarter about the general quarterly average (45,) which is an increase of 73 per cent. on the corresponding quarter of last year, when there was an entire absence of every Epidemic. The deaths from Epidemics this quarter are Measles 6, Scarlatina 5, Small Pox after previous vaccination 2, and Small Pox without previous vaccination 3, Consumption 7 cases.
_		No prevailing disease in the district; there have been a few cases of Small Pox, Measles, Scarlatina, and Chicken Pox, with a few cases of severe Rheumatic Gout.
HEREFORD		Scarlatina has been very prevalent in my district, 12 died in the above quarter.  Measles has prevailed in my district this quarter very much, but only one person has
SHREWSBURY		died of the complaint.  We have had in this quarter a few cases of Small Pox, which I consider is chiefly
_		owing to the inattention of the parents to Vaccination.
	St. Mary seconds private	Deaths this quarter rather above the average; Small Pox has been prevalent in my district, fatal cases 11, every one "without previous vaccination." Not one fatal case has yet occurred "with previous vaccination." The disease is on the decline. The other cases of epidemic or contagious diseases are Typhus 5, Measles 3, Hooping Cough 2, total deaths 85, last quarter 100.
KIDDERMINSTER	Bewdley	The number of deaths this quarter are more than double those of the corresponding quarter last year, Typhus has been prevailing in the district, but few deaths have occurred from it, is not sufficient to account for the increase.
DUDLEY	Dudley	Deaths registered in my district this quarter 168, being 21 less than the last, out of which there are Consumption 31, BowelComplaint 15, and Typhus 21.

sup. reg. district  WALSALL	REG. DISTRICT.  The Borough	There i	s an in		of 16	upon	last q	uarter	, 14 h	ave di	ed_of	Consu	nption	the co	mplain
	Darlaston	The nur	^	f deat											l rathe
WOLVERHAMPTON.	Wolverhampton	The nu Coug	mber h has l	of dea	aths th	is qua	arter hildre	is slig n, 18 d	htly l leaths	elow from t	the ay	verage sease a	altho	ugh H	Toopin
			Scarlat	ina.											
BIRMINGHAM	St. George	The nu pondi Scarl	mber of ing quantina,	narter	of las	st yes	ır, 33	have	died	of Co	onsum	ption,	14 of	in the Meas	corres les, 1
	St. Paul	The nur	mber o	f deat	hs thi	s quar	ter h	as beer	a lit	tle abo	ve the	aver	age, the	ere hav an aggi	ve beer ravate
	Lady Wood	viz. I	mber o umber Fever ase in	of Fe 18, M	ver an leasles	d othe	r Epic Ioopin	demic o g Cou	cases l gh 1,	as be withou	en gre	ater in	propo	rtion t	hereto
ASTON	Erdington	The de		(43) d	luring	the c	narte	r are	above		verag	e of f	ormer	retur	ns, the
LEICESTER	East Leicester		s a dec the dis- juarter	trict I	consi	der ve	ery he	althy,	no epi	idemic	. The	princ			
	West Leicester		mber o										ter; n	ny dist	trict i
LINCOLN		<ul> <li>There has been a marked decrease in the number of deaths this quarter, a decrease of one-third. Last quarter the number was 123, this quarter 92, the difference being 31 A reason may be assigned for it. At this season of the year, there are not such extensive deposits of filth in the city, as there are during the winter quarter, consequently there is less miasma, this I infer from there not being a single case of death I Typhus this quarter. An analysis of this quarter's return will corroborate my forme opinion "that the want of drainage in Lincoln has generated disease, the effects of which will ere long be seen." Out of 92 deaths in this return, 30 have died of Consumption, of which number, only I had reached 50 years, 3 from 40 to 50, the remainder under 40 years. The excess of deaths of males is 22 in 92. The deaths in the public Institutions have greatly lessened.</li> <li>Although the number of deaths is below the average, Typhus has been more prevalent.</li> </ul>													
	North East Park														
	Bulwell	regist	tered o	f Cons	sumpti	on.									
	Ilkeston	The nur the pr lungs instan		of deat ace of the o ter fou	ths thi an Ep rgans ir or fi	s quai idemic princi ve da	rter is during pally ys illn	consider the affected	derably month	y abov of Ap	re the ril, an	averag nong o rminat	ge, ma hildrer ing fat	inly ov in wh ally in	wing to nich the many
.,,,,,,,,			total is	not a	bove t	heave	rage.							Ĭ	1
		regul	over ti ar avei	he last age.	quart	er in t	he nui	nber.r	egiste	red, a	nd of	about	one-thi	rd abo	ove the
STOCKPORT		Seventy Decer regist passin respe	v-four mber of tered 8 ng of ective s	below quarter cases the V	rs. A , a gr accina 28 ve	mall eater ration Lars, 10	Pox 1 number $Act;$ $6, 4\frac{1}{4},$	ias be r than 5 deat	the w	articul hole re thout	arly f egister vaccin	atal thed ed before ation	is qua ore by and 3	rter, l me si	I have
•••••	Stockport (Second) Hyde	Consider Below						not ha	d so f	ew des	iths in		arter s	ince ti	he Aa
	Heaton Norris	From t	into o	peratio	n.										
		I hav last 4 rathe just a Dece gener princ Diarr	e been quart er more averagember, rally, in pal drhæa, sthma	ers; a ethan e; and and ; in a h eaths 3; Pn	ed to a not find find find find find find find find	make I the r cent resent s than state Typhu ia, 4;	a calc Births Fro one, ( Sept Ho is, 3; Teeth with	ulation to be om this 58, bei ember wever, Scarl ning, 3	ofrom 591, the sit wo ng 18, it is , Feve atina, ; Hydoner's	the birde deat uld appless the a cherr has 2; Marocep Inques	rths ar hs 264 pear t an tho sering been s leasles balus,	nd dead that 66 se of 1 fact 1 somew 5, 2;	hs reg is as : death March, hat th hat pre Hoopin	istered 2.238 to s wou 40 les e distri- valent ng Cor	l in the to 1, or to be a ss than rict is The ugh 1
		1					1				92	- 00	1 00	v2	1 10
		Under 1 Year	2 Years	3 Years	l Years	5 Years	10 Years	15 Years	20 Years	30 Years	40 Years	50 Years	60 Years	70 Years	80 Years
		17	5	4	3	0	5	0.	3	5	8	5	4	3	6
	Hazelgrove	There a												v the a	verage
MACCLESFIELD		numb rema	ber. Trkably	Che m health	edical y.	man	of this	villag	re has	just :	inform	ed me	that t	be dis	trict is
	, and the second	preva of th	alent, d e Lung	of whice gs, and	h 1 pe	erson h Scarla	as die tina.	d, 5 of	Inflan	matio	n of the	e Ches	t, 4 of	Inflam	matio
	Rainow	Above The nu prece		of dea	aths re	egister	ed du	ring t	his go	arter	is 44,	being	8 les	s than	in the

REG. DISTRICT. SUP. REG. DISTRICT. ACCLESFIELD .... Prestbury ............ Rather above the average, but no particular cause to be assigned for it. Influenza has been prevalent, but in no case, strictly speaking, fatal; in one case only of Pneumonia death was accelerated by it. Out of the 25 deaths registered by me this quarter, one-sifth have been from Phthisis. REAT BOUGHTON. Hawarden..... The deaths are 16 below the average. The number of deaths quarter ending September, 1843, was...

Do. do. December, , ...

Do. do. March, 1844, ... 39 21 128 83 June, 62 299 97 16.839 Number of Deaths Number of Deaths Diseases Diseases City County City County Total Total part part part part 276 Small Pox Brought forward 204 Teething Worms 2 2 2 Scarlatina 4 2 Ulceration Disease of Stomach 11 9 Hooping Cough 78 Croup Diarrhœa . . Jaundice
Disease of Liver
Ditto Kidneys
Gravel  $\frac{2}{2}$ 2 2 •• Cholera 2 Influenza 3 Remittent Fever 4 2 Typhus
Hydrocephalus
Apoplexy
Paralysis
Convulsions
Cattarrh 11 3 50 14 4 61 Dropsy Rheumatism 5 3 6 Decay of Nature Coroner's Cases Hœmorrhage 11 7 11 18 61 15 76 13 Cancer Disease of Spinal 3 1 2 ì Hydrothorax Asthma Marrow 1 2 i Scrofula Mortification Consumption & Decline Premature Birth Disease of Heart Inflammation of Lungs 55 28 83 1 1 i Purples 3 23 8 31 Bowels, &c. Carried forward 204 72 276 Total..... 299 97 396 IVERPOOL ...... Great Howard Street .. I have registered 20 deaths from Small Pox, 18 not vaccinated, and 2 after vaccination one of which was vaccinated two days after it was taken ill. ..... Dale Street ........ The deaths in my district are fewer by 83 in the general number than the previous quarter, the district I conceive being in a very healthy condition, still there is an increase of 10 deaths more than the last quarter in the disease of Small Pox, without previous vaccination. The deaths this quarter are 80 less than the last, and 25 below the Spring quarter of 1843; this is the smallest quarter since the commencement of the Registration Act, which has now completed its seventh year; the number for the whole year is 40 less than the preceding year, and 57 under the year ending June, 1842. The principal diseases are Hydrocephalus 14, Phthisis 37, Bronchitis 16, Pneumonia 22, Diarrheea &c. 13, Hooping Cough 9, Teething 9, Dropsy 8, Croup 5, Measles 6, Inquests 17, Delirium Tremens 2. .. St. Thomas There is a decrease in my district of 75 deaths, as compared with the corresponding quarter of last year. ... Islington ..... ..... Saint George .... The number of deaths in my district this quarter is 42 less than the corresponding quarter of last year, and 71 less than the preceding quarter, consequently the district is remarkably healthy. There is an increase of 27 in the number of deaths as compared with my last quarterly report; particulars—Teething 2, Epilepsy 1, Small Pox 3, Scalding 1, Decline 12, Convulsions 15, Hæmorrhage from the Navel 1, Dropsy 6, Caneer of the Womb 2, Water on the Brain 3, Consumption 12, Tubercular Phthisis 1, Decay of Nature 7, Palpitation of the Heart 3, Scarlatina 6, Inflammation of the Lungs 5, Paralysis 2, Child Birth 1, Asthma 2, Apoplexy 1, Measles 1, Croup 5, Liver Complaint 3, Hæmatemesis I, Hooping Cough 3, Inflammation of the Chest 2, Inflammation of the Brain 2, Inflammation of the Bowels 2, Ulceration of the Stomach 1, Hydrocephalus Acutus 1, Inquests 3. WEST DERBY ..... West Derby .... .. Great Crosby ..... Little or no Epidemic disease this quarter.

The deaths this quarter are below the average; no particular Epidemic prevailing; my
district is in a very healthy state except for Consumption, 70 persons out of 240 have
died of that complaint.

 The number of deaths in my district is above the average of former quarters, owing to the prevalence of Hooping Cough or what is generally called Chin Cough.

· · · · The deaths in this district are about the average.

..... Mellor

Oswaldwistle ...

Witton

The number of deaths this quarter is only one more than in the preceding quarter, although two epidemic diseases have been prevailing in my district, viz.—Small Pox and Hooping Cough.

SUP.REG. DISTRICT. REG. DISTRICT.	
	. There are 30 deaths less this quarter than in the corresponding one of last year; the
	prevailing disease, Consumption.
Longton	Of the 29 deaths, 9 are of children under the age of 1 year. The causes of death are Phthisis 7, Debility from birth 6, Scarlatina 2, Typhus 1, Enteritis 1, Convulsions 1, Dropsy 2, Ileus 1, Gravel 1, Diabetes 1, Meningitis 1, Serofula 1, Old age 1, Sudden Death 1, Coroner's Inquests 2.
Walton-le-dale	The number 35, is the same as the June quarter for 1843, but 8 below an average of the corresponding quarters since 1837.
ROCHDALE Castleton	The number of deaths in my district is above the average owing to the prevalence of Small Pox.
Wardleworth	<ul> <li>Small Pox is still prevalant in my district; having had 17 cases without previous vaccination, and two attacks after vaccination.</li> </ul>
BURY Bury, South	The return of the number of deaths registered during the last quarter shews an increase of rather more than 3 deaths over the average of the 5 previous corresponding quarters, but if the increase of population be taken into account it will be under the average.
	The deaths are as follows:—Measles 9; 8 before the 4th of May and 1 in June. Diarrhoea 4, Scarlatina 5, all before the 2nd May; Premature Births, 4; 1 lived 5 minutes, 1, 3 hours; 1, 14 hours; and 1, 2 days. Typhus 1, Abortion with Syncope 1, Childbirth 1, cause not stated; Inquests 2. (1 Cutting his throat, 1 Apoplexy). My district is in a very healthy state at present.
	<ul> <li>There have been more deaths from Measles in this quarter than usual; and more cases of Typhus than in former quarters. There have been a few deaths from Small Pox in persons not vaccinated but on the whole my district is in a very healthy state at present.</li> </ul>
Tottington	Measles is not so prevalent as it was.  Rather below an average for the corresponding quarters; my district is in a healthy state
BOLTON Little Bolton	at present and no epidemic prevailing.  This district is more healthy than it has been for many previous months, and there has the property of deaths. No particular discrete has
Farnworth	consequently been a decrease in the number of deaths. No particular disorder has been prevalent during the last quarter.  The number of deaths this quarter is two above the general average; but fewer by 28
	than in the corresponding quarter of 1843. There has been no epidemic and the district is considered healthy.
WIGAN Wigan	The number of deaths, although decreased, still continues above the average in Wigan; 29 have been registered from Phthisis or Consumption, 18 from Scarlatina, and 42 from Small Pox; only one of the 42 persons had been vaccinated.
Pemberton	. 65 deaths being 15 above the average. 25 cases of Small Pox — 10 males, 15 females; not 1 of which had been vaccinated. 13 cases of Consumption, 6 Decay of Nature, Convulsion Fits 5, Dropsy 2, Asthma 1, Brain Fever 1, Inflammation of the Chest 1, Inflammation of the Bowels 1, Paralysis 1, Typhus Fever 1, Teething 1, Debility 1, Premature Birth 1, Apoplexy 1, Burning 1, Kick from a horse 1, Coal Pit Accidents 2.
Ashton	<ul> <li>My district is in a very healthy state; the present return is rather below the average of former Spring quarters. The most remarkable feature on the face of the register is the mortality of aged persons. More than one-third of the 25 deaths registered consists of persons whose average ages reach 75 years.</li> </ul>
PRESCOTT St. Helens	68 Deaths. This district has been particularly healthy this quarter; 54 below the average of the corresponding quarters of 1841-42-43.
····· Prescot	<ul> <li>This district has been in a very healthy state during the last quarter, and the number of Deaths has been considerably below the general average, 13 have died between the ages of 60 and 70 years, 7 between 70 and 80 years, and 1, 84 years old.</li> </ul>
	There is no particular disease prevalent in my district this quarter.
	There have been only 148 deaths registered during the last quarter; which number is about 30 less than the average. The unusual healthfulness has been owing partly to the more comfortable condition of the people in consequence of improved trade, partly to the remarkable fineness of the weather.
MANCHESTER Ancoats	Total deaths 266 is much below the average—102 died under 2 years of age, of Scarlatina 1, Measles 12, Inflammation of Lungs 18, Inflammation of Bowels 2, Consumption 45, Fever 3, Croup 5, Child Birth 1, Dropsy 8, Convulsions within 1st Month 23, Ditto within 3 Months 8, Ditto 1 year or upwards 18, Teething 14, other Causes 108.
St. George	The number of deaths registered this quarter is considerably below the average. The principal causes have been Consumption 33, Pneumonitis 22, Convulsions 20, Dentition 13, Asthma 12, Hooping Cough 4, Hydrocephalus 6, Diarrhœa 6, Measles 6, Tabes Mesenterica 5, Bronchitts 3.
	257 deaths.—Independently of the Workhouse, 1nfirmary and Coroners' Inquests, the deaths in my district in the last quarter are 111; being 51 less than the corresponding quarter of 1843. This decrease is owing to epidemic diseases having but little prevailed, and to the improved condition of the population.
	Deaths for the quarter under the average.
Cheetham	The deaths are much below the average.  Deaths 57.—No epidemic disease; but I find that in the June quarters of 1842, 1843, and 1844, the numbers are greater than in any other quarters. I know of no particular cause, unless it may be attributed to the great variation in the weather, viz. dry hot days, and frequently cold nights.
	The number of deaths registered during the last quarter is below the average, although the number of deaths in the Infantry Barracks has been, I believe, greater than in any previous quarter since the Registration Act came into operation. In Barracks 10 deaths—1 serjeant, I corporal, 8 privates.
···, ···· Greengate: › · · · · · · ·	About 40 below the quarterly average and about 20 below the average of the June quarter, although an unusual number of accidental deaths has taken place. Number of deaths in April 68, in May 87, in June 51.
ASHTON Ashton Town	Less than the average of corresponding quarters since commencement of Registration Of the 135 deaths 48 were without medical attendance, 76 were attended by qualified practitioners, and 3 by druggists.
	The number of deaths in my district is about the average of the two preceding corresponding quarters. There is no epidemic prevalent in this district at present.
Denten	Much below the usual average.
Dukinfield	My district is free from any epidemic.  There are 11 deaths less this quarter than in the June Quarter of 1843.
	And are are decline less time quarter than in the June Quarter of 1045.

CHD DAG Prompto	DPG DIGMOSOM	
SUP. REG. DISTRICT.  ASHTON	Mottram	The number of deaths is much below the average. My district is very healthy.
	Middleton	The number of deaths this quarter is still under the average; my district is in a very healthy state.
	Royton	Attended by qualified persons, 29; without medical attendance, 9; 38 deaths: being 5 less than the average of the three previous corresponding quarters. This district has been more than usually healthy this quarter.
	Chadderton	The causes of death this quarter are of an ordinary character; there are five Coroners' Inquests.
SHEFFIELD	Park	An average number this quarter.
	Sheffield South	63 deaths. The number registered is less than usual, being only 28; the weather being fine and cool during nearly the whole time.
		My district is in a healthy state, the deaths being 55 less than last quarter.
		7 deaths. It appears that a less number of deaths have occurred during this quarter than in any of the previous corresponding quarters, which have been respectively 10, 16, 9, 13, 18, 19.
HUDDERSFIELD		There have been 2 cases of Typhus, and 2 cases of Inflammatory Fever within the quarter. Consumption appears to have been prevalent; 14 deaths are referred to that disease.
	Honley	The number of deaths registered during the last quarter is about the average. No epidemic has prevailed to any considerable extent. There have been 2 cases of Typhus, 1 of Influenza, and 5 of Consumption.
	Lockwood	The number of deaths in my district this quarter is somewhat above the average of the corresponding quarters of the years 1840, 1841, 1842, and 1843. No less than five-ninths of the present return being children under 4 years of age; the prevailing diseases were Hydrocephalus, Convulsions, Pneumonitis, Teething, &c. likewise 3 of Hooping Cough, 2 of Premature Birth, and 1 of Thrush.
		The number of deaths registered in this district during the quarter has been rather above the average; there has been no epidemic disease.
HALIFAX		The number of deaths (131) this quarter is 25 below the average of the corresponding quarters of the 6 preceding years, and is 54 below the last June quarter of 1843.
	Brighouse	Above an average; the increase appears to be with old people.
	Rastrick	The number of deaths (19) is under the average.
	Ripponden	The number of deaths this quarter is 42, being 12 more than in the corresponding quarter of last year. This is attributable to the prevalence of Scarlatina and Typhus.
RR4DFORD	Northowram	The number of deaths in this quarter exceeds those in the corresponding quarter of 1842 by 14; arising chiefly from a rapid increase of population in this district.
	East-end	I have an Increase in deaths this quarter of 19, as compared with the quarter ending March, which 1 consider to have arisen from Small Pox and Meales. Deaths from Small Pox.—Males: 1, 34 years; 1, 10 years; 1, 5 years; 3, 3 years; 2, 2 years; 1, 1 year; 1, 6 months; 1, 7 mouths; 2, 8 months. Females: 1, 18 years; 1, 10 years; 1, 7 years; 2, 4 years; 1, 3 years; 2, 2 years; 1, 6 months; 1, 10 months. Males, 10 years; 1, 3 years; 2, 2 years; 1, 6 months; 1, 10 months. Males, 10 years; 1, 3 years; 2, 2 years; 1, 6 months; 1, 10 months.
		vaccinated. Measles.—Males, 2; Females, 3; total 5.
	West-end	Deaths 164. I have 40 deaths this quarter more than the average number, arising chiefly from the Small Pox and Convulsions among children.
	Cleckheaton Thornton	The number of deaths is below the average and the district is very healthy.  The number of deaths, (75) though there has been no epidemic disease prevailing in
		my district is about 25 per cent. above the average. During the cold and dry northeasterly winds in May, the mortality was very great, especially among aged people; and more deaths took place in that month than in the other 2 months in the quarter.
86866666		About an average; 1 case of Typhus, 2 of Croup.
01000000		Scarlatina and the Croup have proved fatal in several cases.  The number of deaths in this district during the last quarter has been above the average,
	Pudsey	which must be attributed to general causes, as no epidemic disease has prevailed, but Pulmonary complaints appear to be on the increase.
	North Bierly	Deaths 60.—The average of the corresponding quarters is 56. (Deaths have varied from 48 to 70 in the quarters alluded to.) I have had one death from Small Pox, not vaccinated, during the quarter.
		The number of deaths are more by 12 this quarter than last; though I do not perceive any particular complaint to be prevalent. Included in the number are 19 cases of Consumption, and 3 cases of Small Pox, (not been vaccinated).
		The number of deaths (205) is precisely the same as was registered in the corresponding quarter of last year, but considerably below the average of former years. The number of deaths of children under 5 years of age is 84, being much fewer than has been registered in former quarters. There has been no excess of deaths from any epidemic or other disease, a considerable number of the deaths registered having been caused by Phthisis, Old Age, Paralysis, Convulsions, &c.
HULL	St. Mary's	The number of deaths this quarter is below the average.
YORK	Micklegate	Below the average; 2 cases of Small Pox, (without vaccination) and 3 of Scarlatma.
	Walmgate	The number of deaths registered this quarter is 115, being 22 more than last, and 40 more than the corresponding quarter of last year. There have been 17 deaths from Small Pox (only 4 of which had been vaccinated) 4 of Scarlatina, and a considerable number of cases of Consumption.
_	***	13 deaths, which are less than the average; a few cases of Small Pox have occurred, but none fatal.
2		In this quarter there are 16 deaths, a greater number than the average; the increase is owing to Scarlatina, which was and still continue very prevalent in one part of my district.
SUNDERLAND		29 deaths. About one half the number registered in any of the corresponding quarters since the commencement of the Registration Act.
04000		The number of deaths during the quarter has been above the average, owing to the prevalence of Measles.
0,000	South	12 above the average. Measles prevailing in my district. I have had 7 cases of Measles many of them terminating in Pneumonia.

SUP. REG. DI	STRICT.	REG. DISTRICT.	
GATESHEAD		Winlaton	Below the average.
TYNEMOUTH		Tynemouth	Deaths 83. The present return gives an excess of 17 above the number registered during the same period of last year, yet the mortality is less, by 2 four-sixths, than the average of the 6 previous Spring quarters. 30 infants under 3 years of age, have died this quarter from Teething, Convulsions, Debility, and Hooping Cough; which latter disease has been very prevalent, and in numerous cases fatal. The deaths of aged persons amount to 18, viz., 4 males and 14 Females, all of whom exceeded 70 years of age.
		North Shields	The number of deaths (75) registered this quarter is about the usual average. Of the 43 Males. 24 are infants and children under 10 years of age, from ordinary causes. Of the 32 females, 9 are infants of tender age. Included in the total number are 7 Inquests, 5 males and 2 females.
		•	The number of deaths is below the average, and there has been little or no sickness in the district.
NEWCASTLE TYNE	UPON	All Saints	Deaths 103. My district is particularly healthy; and the present Return is at least 30 below the average.
CARLISLE		St. Mary	The number of deaths (89) this quarter is about an average return, but one-third below the last quarter. The increase last quarter was from <i>Measles</i> and <i>Hooping Cough</i> which have now abated.
		St. Cuthbert	57 is an average number of deaths; there does not appear to be any epidemic prevailing in my district. There have been 2 deaths from <i>Measles</i> and <i>Hooping Cough</i> , and 2 Inquests, all the rest are chronic diseases.
		Stanwix	Deaths 21. I have 12 more deaths this quarter than in the corresponding quarter of 1843. The only causes of the increase are <i>Hooping Cough</i> and Water in the Head only 1 case of <i>Measles</i> this quarter.
		Buryh	Deaths 10. The increase of deaths in my district for the present quarter above the average number, is from accident not from any epidemic prevailing at the present time.
	• • • • • •	Dalston	The deaths in this district have been 10; and there are 9 deaths fewer than in the corresponding quarter last year.
6 0 0	• • • • •	Wetheral	This is below the average number, it being one-fifth less than in the corresponding quarter of last year.
			The number of deaths registered this quarter is rather above an average, still there is no particular disease in my district; out of the 30 entries 13 are children under 2 years, and 6 from old age, namely, 74, 75, 87, 83, 70, 74.
		Grayrigg	
		Milnthorpe	This district is very healthy, and the number of deaths is the same as in the quarter ending 30 June, 1843.
			My district at this time is in a healthy state.
MERTHYR TY	DFIL .	Merthyr Tydfil, Upper	The number (244) is considerably above the corresponding quarter of last year, being more than double; resulting from Small Pox, Scarlatina, and epidemic fevers.
		Lower	The deaths are still considerably above the average, there are 16 of Scarlatina and 31 of Small Pox in this quarter.
NEWTOWN		Tregynon	The number of deaths this quarter is the same as in the previous quarter, and I consider the district to be in a healthy state.
			The number of deaths in this quarter is rather above the average of corresponding quarters. Fever is prevailing much in my district, of which 16 have died, 7 Bronchitis, 14 Phthisis, the other causes are similar to last quarter.
HOLYWELL		Holywell	Total deaths, 79. Deaths of children under 4 years, 41; Measles 11, out of that number other causes, 30; therefore, this disease is the most prevalent. Total deaths less in this than the March quarter 20, but 15 more females than males; the preponderance of deaths being greater in the males than females in the preceding quarter.

GENERAL REGISTER OFFICE, JULY 25th, 1844.





# A QUARTERLY TABLE

OF THE

# Mortality

IN A

# HUNDRED AND FIFTEEN OF THE DISTRICTS

OF

# ENGLAND

(INCLUDING THE PRINCIPAL TOWNS,)

SHEWING

THE NUMBER OF DEATHS REGISTERED IN THE SIX YEARS 1838—43, THE AVERAGE NUMBER OF DEATHS IN THE FIVE SUMMERS 1838—1842;

AND THE NUMBER OF DEATHS IN THE SUMMER QUARTER 1844, ENDING SEPTEMBER 30TH.

PUBLISHED BY AUTHORITY OF THE REGISTRAR GENERAL.



LONDON:

HARTNELL. PRINTER, RED LION COURT.

1.381

### was to be a stress to the

A service of the servic

en de la companya de

# THE STREET PRODUCTION OF THE STREET

# Carlotte State State Control of the 
transfer to the second of the

Teta 2

### TABLE

OF THE

### MORTALITY

In 115 of the Districts of England (including the principal Towns): shewing the Number of Deaths Registered in the 6 years, the average Number of Deaths in the 5 Summers, 1838-42, and the Number of Deaths in the Summer Quarter of 1844, ending Sept. 30th.

					-			1838	-42	ummer ling 1844						1			1838	-42	mmer ing 1844
FRICTS	Popu-	De	aths 1	Registe	red in	the ye	ears	Qna • Av	rterly erage	the S r, enc 10th,	DISTRICTS	Popu-	I	eaths 1	Register	red in t	he year	3.	Quar • Ave	rage	in the Summrter, ending 30th, 1844
	1841	1838	1839	1840	1841	1842	1843	Of Five Years	Of Five Sam- mers	Deaths in Quarte Sept. 3		1841	1838	1839	1840	1841	1842	1843	Of Five Years	Of Five Sum- mers	Deaths in Quart Sept.
ropolis											N. Midland division										
Districts	301326	8114	6538	1 1	6599	6826		1751	1617	1822	Leicester	50932	1170	1291	1730 817	1358	720	1255 786	<b>35</b> 0 184	312 159	339 143
Districts	366303	8944	7890	9108	9333	8550 8873	9038 9547	2112 2395	1966 2238	2293 2190	Nottingham	36110 53080	1209	731 1540	1794	1239	1429	1393	361	366	391
al Districts Districts	374759 393247		9751 10359	10063	- 1		10960	2640	2567	2547	Basford	59634	1045	1200	1549	1163	1245	1216	310	266	233
Districts	479469		11562	12329	11872	11918	12575	2850	<b>2</b> 632	2973	Derby	35015	769	832	926	914	909	922	217	196	200
Fotal	1915104	53546	46100	47030	46069	46114	49332	11749	11020	11825	N. Western division	-									
astern								1			Stockport	85672	2410	2591	2574	2048	2073	2203	585	505	462
tone	35210	737	726	700	648	. 815	774	181	161	160	Macclesfield	56018	1315	1528	1745	1419	1403	1551	370	307	283
ton	46742	916	898		1102	1126	1117	251	254	233	Gt.Broughton, inc. Chester	49085	1040	1237	1384	1246	1189	1091	305	253	208
Wight	42547 53036	655	630 1180	1 1	891 1211	721 1152	703	182	170 272	187 345	Liverpool	223054	6627	7435	8470	7556	7407	7458	1875	1830	1945
ea Island	23044	1264	479	1 1	416	469	471	116	117	100	West Derby, \ (adj. Liverpool)	88652	1625	1746	2206	2202	2405	2312	509	498	602
sor	20502	381	367	402	395	394	314	97	83	99	Blackburn	75091	1612	2190	2140	1692	1665	2031	465	381	474
idland											Preston	77189 60577	1726	1812	2637 1868	2026	2050	1938	513 396	424 325	450 316
bans	17051	378	310	331	300	340	291	83	74	100	Bury	77496	1901	1996	2170	2064	1987	1832	506	405	380
mbe	34150	788		1 000	751	679	748	187	163	180	Bolton	97519	2432	2857	2900	2572	2607	2596	668	559	534
d	19701	359		406	450	414		102	96	90 1 <b>62</b>	Wigan	66032	1483	1912	2144	1561 997	1588	1832 920	434 260	357 213	353 174
ampton	28103 31767	689 676	687	670 743	693 714	622	677 762	168 171	157 146	175	Prescott	43739 93736	916 2140	1056	2399	2431	2427	2529	591	508	653
ridge	24453	563		639	659	465	597	144	134	133	Manchester	192408	6706	6774	6489	5821	6151	6263	1597	1462	1441
stern											Salford	70228	2207	2449	1998	1971	1977	2098 4391	530 1196	501	417 918
ision	18900	#00	437	488	381	417	393	110	106	124	Ashton	173964	4488	5489	4873	4282	4786	4091	1150	1000	210
ch	17790 25254	533 694	630	593	499	594	551	113 150	144	135	York division							1			
ich	61846	1373	1852	1	1362	1582	1295	396	22-	306	Sheffield	85076	2429	2703	2410	2231	2182	2169	598	530	494
outh	24031	417	440	504	493	516	510	118	117	115	Huddersfield	107140 109175	1847 2098	2080	2243	2122	1996 2272	2451 2459	514 566	460 474	447
estern											Bradford	132164	3093	3251	3176	3041	3323	3344	794	728	861
es	22130	426	510	481	429			115	99	84	Leeds	168667	4289	4388	4488	4360	4615	4329	1107	1042	997
ester	23380	1			456 920			107	300	1	Hull	41130	1060	1023	1209	1206 1039	1042	1128	289 274	284 243	247 247
r	31333 47105	1		1				102	1		York	47779	876	1217	1200	1000	1100	1010			
onth	36527	1		1			3	1	201	257	Northern division								- 0.0		
uth		1	1	1 000		1	1	3	1 000		Sunderland	56226	1479	1524	1513	1512	1357	1247	<b>3</b> 69 <b>2</b> 38	353 229	267 228
ance	69232	į.	į.	-00		1			1		Gateshead	38747 55625	881	968 1285	1016 1243	989 1340	909	984 1229	315	296	242
	00202	100	100			1000					Newcastle-on-	71850	1829	2117	1957	2104	1726	1922	487	485	414
estern vision		1:									Tyne	36084	855	739	901	925	840	762	213	162	165
ol	00000		2 167	****	1	1510						35676	784	718	696	674	597	690	173	152	146
on		1	1	-		3					77 3 - 1	34694	729	825	821	601	714	666	184	162	154
teuham	1			1 000	1	1			196	191	Welsh										
ford				1							41	50834	1150	1206	1374	1458	1109	1116	315	260	244
wsbury	1	1			1	1		1	3 00		Pontypool	25037	561	615	<b>5</b> 63	648	545	514	147	123	97
cester lerminster	2940					1	}				Merthyr Tydfil	52864	1516	1246	1566	1423	_	1183	343 129	283 114	397 101
ley	0000	1	1	2025	1			000			f	25958 39542		468 823	619 966	<b>53</b> 9	420 809	477 917	217	185	159
sall		1	- 1		1		1	1			1 vr			791	864	800	1	815	204	162	177
verhamptor stanton'		1 1	- 1	- 1	1	3 2319 7 96		1 - 00				38105	ī	589	632	597	624	575	154	126	122
ningham	1	1				3 3579		90	1 87				109321	116505	124664	114664	115834	113869	29049	26063	26959
II'	5092			3.	1	1		1 20			4				-	_	_				
entry	3102	8 68	72	61	6 90	5 89	0 83	8 19	1 17	6 231	GRAND TOTAL	6578912	102867	162605	1/1094	100733	101948	100201	40/98	37083	30/84
			1			1			1	1							1	1	1		-

<sup>\*</sup> The Deaths in the Metropolis for the Years 1840 1-2, have been derived from the Weekly Tables, the Returns for each Year comprising 52 Weeks, or 364 Days. The last quarter in the Metropolis ended Sept. 28, 1844. The Returns from other Places are for the years ending December 31, and the quarter ending Sept. 30th.

<sup>\*</sup> Wandsworth District is included in the return for the Metropolis.

### A TABLE OF THE MORTALITY

IN THE

### Metropolis,

### SHEWING THE NUMBER OF DEATHS FROM ALL CAUSES,

Registered in the 13 Weeks ending Sept. 28th, 1844.

CAUSES OF DEATH	Quarterly Average of Deaths in the 5 Summers of 1839-43	Deaths in the Summer Quarter ending Sept. 28th 1844	CAUSES OF DEATH	Quarterly Average of Deaths in the 5 Summers of 1839-43	Deaths in the Summer Quarter ending Sept. 28th 1844
ALL CAUSES	11131	11825	IVLaryngitis	6	7
SPECIFIED CAUSES	11083	11797	Quinsey	21.	32
I Zymotic (or Epidemic, Ende-	23 <b>57</b>	3243	Bronchitis	73:	140
mic, and Contagious) Diseases	2007	0240	Pleurisy	15	23
SPORADIC DISEASES:-		'	Pneumonia	656	617
IIDropsy, Cancer, and other Dis-			Hydrothorax	41 113	38
eases of uncertain or variable Seat	1312	1239	Phthisis or Consumption	1833	1681
III.—Diseases of the Brain, Spinal	7000	2000	Disease of Lungs, &c	126	140
Marrow, Nerves and Senses	1968	1929	V.—Pericarditis	8	28
IV.—Diseases of the Lungs and of the	2884	2782	Aneurism	8	8
other Organs of Respiration ( V.—Diseases of the Heart and Blood )			Disease of Heart, &c	218	290
Vessels	235	326	VI.—Teething	301	233
VI.—Diseases of the Stomach, Liver,	1085	1027	Gastritis		15
and other Organs of Digestion			Enteritis	3	264
VII.—Diseases of the Kidneys, &c	63	90	Peritonitis		26
VIII.—Childbirth, Diseases of the Uterus, &c	105	124	Tabes Mesenterica	84	136
IX.—Rheumatism, Diseases of the			Ascites	6	6 27
Bones, Joints, &c	65	84	Ulceration (of Intestines, &c.).	23	18
X.—Diseases of the Skin, Cellular	15	12	Hernia	17	20
Tissue, &c			Colic or Ileus	26	37
XII.—Violence, Privation, and In	686	648	Intussusception	9	3
temperance	308	293	Stricture	7	5
I.—Small Pox	130	556	Disease of Stomach, &c.,,	6 66	13 86
Measles	404 476	255 1020	Disease of Pancreas	.4	4.0
Hooping Cough	275	167	Hepatitis	17	21
Croup Thrush.	80 111	76 120	Jaundice	29	28
Diarrhœa	331	414	Disease of Liver, &c	113	89
Dysentery	59 54	44	Disease of Spleen	3	1.40-9
Influenza	12	. 8	VII.—Nephritis	4 2	5
Ague	3	6	Diabetes	4	2
Typhus	352	424	Cystitis	3	2
Erysipelas	55 9	··· 85	Stone	8	7
Syphilis	1		Stricture	5	17
II.—Inflammation	79	7	Disease of Kidneys, &c,	36	56
Hæmorrhage	37 377	30 312	VIII.—Childbirth	71	82
Abscess	41	. 30	ParameniaOvarium Dropsy	2	1
Mortification	55 5	45	Disease of Uterus, &c.	5 27	38
Scrofula	30	45	IX.—Arthritis	.6	
CancerTumor.	111 23	121 11	Rheumatism	30	1.
Gout	10	10	Disease of Joints, &c	35	52
Atrophy Debility	124 258	201 287	XCarbuncle	1	1
Malformations	13	25	Phlegmon	.4	1
Sudden Deaths	150	108	. Ulcer	- 4	1
III.—Cephalitis	156 475	164 413	Fistula	4	, 3
Apoplexy	197	237	Disease of Skin, &c	5	6
Paralysis	168 781	166	XI.—Old Age	686	648
Tetanus	4	721	XII.—Intemperance	11	8
Chorea Epilepsy	.4	. 2	Privation	5	3
Insanity	41 11	54 19	Violent Deaths	292	282
Delirium Tremens Disease of Brain, &c	23	22	Causes not specified	47	28
Division of Diam, the control of the	112	122			
				-	

P
口
BL
A
TAB
ICAL
7
25
OLOGI
H
0
2
0
EOR
-
MET
$\rightarrow$
$\Box$
RI
AR
A
D
0

Compiled from the Weekly Tables furnished to the Registrar General by the Astronomer Royal.

Deaths registered in the Metropolis from Snall Pox, Measles, Scarlatina, Hoping Cough, Typins, Diarrina, Dysentery and Cholera, Influenza, Consumption, and other diseases of the Langs; the numbers at each age, and the total deaths ( except violent and sudden) in each of the 13 weeks ending Sep. 28, 1844.

	10	eviau	esths)	eaths from all causes violent and sudden de	α	843	815	867	1034	973	816	906	903	763	868	810	848	862	11435
	90	len	1	sparadn pue	09	661	127	133	162	143	149	137	9	57	e)	01	000	-	
	Three	sudd			-								146	127	153	132	128	137	
	aths at	violent and sudden Deaths)		09 93 9	-	366	253	261	98	380	287	281	267	310	239	255	249	273	3367
	Arei	viole		61 03	0	437	434	473	575	549	592	488	490	426	200	400	471	451	6248
	ï		sSun	f and lo easses the l	0	101	200	19	56	95	79	98	3,8	98	18	76	8	96	011
			uo	hthisis or Consumpti	d	137	134	129	130	145	132	130	116	116	129	121	124	138	1691
-				tzuənyn		C1	66	2	-	2				•	pm	eo			8
					_	13	proj	45		-		20	92		7	4	*E		10
		101688	and Ci	iarrhaa, Dysentery,	_			*	57	17	7.1	· · ·	es	 	27	24		25	505
				snydá	L	44	28	34	37	33	89	26	3	31	39	=	33		424
				leooping Cough	H	17	12	13	S	13	13	00	01	12	71	16	14	11	167
				enitalina	s	58	69	72	22	75	80	89	63	85	86.	2,0	87	82	1020
The state of the s				leasies	u	21	61	24	55	13	14	5	53	21	56	· 82	80	15	258
				nsil Pox		36	43	37	65	4.4	46	88	47	36	49	37	27	37	\$56
1	-					_		-	4								-		
				ayab () eshoni ni nini	H	1,28	0.71	0.07	:	0.70	0.44	0.92			0.19	0.46	5 0.59	0.02	5-38
			01-0 °F	lean amount of Cloud	v	8.7	8.3	5.4		6.9	5.0	7.7	7.1	4.2	5.6	8.00	6.5	3,9	6,3
	ıttət	nəvən	ontal n	he amount of Horiz of the air in each we	L	miles 450	920	1115	582	1950	1142	1315	PF 00	485	710	520	830	480	67.0
		all all	1 1	dean for the week	-	0.01	94	0,5	0.04	1.1	0.7	2.0	0.24	0.03	0.1	0.04	0.2	90°0	0.29
	( D	ho on	foot	тре меек	-	0.0	0.8	7.0	1.5	8,5	7.0	13.	3:5	3.0	3.0	0.1	0.4	3.0	0.00
	WIN	I wi out	square	i sansesure i										rai		· X	* :		
		Proces	CO .	notheral direction	P	N.E.	N.N.W.	W	N.E.	W.S.W.	S.W.	W.	W.	N.W.&	E.N.E.	and W.N.W.	W.S. Pand N.E.	N.N.E.	
	110	Naa.u	SHIDS SHIDS	an average of 25 yea		40	ıū	4		0	0	64	62	F-	zņ.	rō.	00	F.	2.0
	npe-	n tem	he mea	ifference between training week, so it is out to state the section of the section	1 0	-3,5	1 1	5.4	+ 5.1	14,9	-4.0	7	. 50	-3.7	13.5	+0.5	9,0	-3.7	18
		are	-1	each day, 6 observe	· ·	0.5	2.0	1,2	1.2	1:0	1,0	0:5	1,2	1.1	The State of the S	8,0	0.1	*	8.0
	the des	nperati	rature	vations dean of the least o	N														
	Differen	oint temperature	tempe	Mean of the greater on each day, 6 observed	0	6,8	14.7	14.4	16.1	14.0	13,5	** **	12.6	11.5	10.7	9,7	7.7	11.4	11.8
-		· Od		someteffice of the series	0	3.9	7.6	6.7	7.00	9'9		63 00	6.3	5.6	eo cri	4.5	2.7	63	
		ring	Lowest on the	dean of 7 obser-	u =	46.2	49.5	40.2	43.8	44.7	42.1	40.8	8 44.4	8 38.1	8 48,0	16.6	0 42.6	83.5	43.5
		Self-Registering		Valions the mook	-	\$ 39.3	94.2 42.5	90.9 31.5	105.3 40.5	89.7 37.7	92,5 34,5	90	86.9 36.1	31.	96.2 36.1	86.2 34.0	64.3 30.0	85.4 30.0	90°0 30°0
	R S	Self-F	Highest in the	Jean of 7 obser-	_	88	101,0 94	9	115.0 108	110.6 86	13	6.7 80	98.86	8.0	104.6	95.4 86	88°3	92.0 8	-
	TE	A.		1		55.0 104	53.9 10	1.2 97	59.7 11	51.8 11	52.8 99	54.4 96	52.1	50.9 100.	58,9 10	58.6	53.8	48.2 9	53.6 115.0
ł	OME	Dew		dean of 72 results		58.9	61.5	7.9 51	1/2	58.8 5]	50.7	55,2	88 8.4 55	56.5	62.7 51	58,1 5	56.5 5,	52.0 48	58.9
1	R M			Difference dean of 7 sobservation	-	12.3	13.6	18,3 57	20.8 67	14.0	16.4	10.7	16.1	19.3 5	15.8	12.8	13.0	18.7	5.5
-	THE	ean	4 Same	the Lowest on each stone from 6 observations	-	62 62	55,0	48.7	57.7	1.19	80.5	52.8	50.4	47.2	55.6	51.0	50.2	43,8	51.4
		M		suomenjasgo 9 moli	-1-0	9.49	9.89	0.70	78.5	65.1	6.99	63.3	66.5	66.8	73.6	64.7	63.2	62.5	6.99
				Lowest during the we	-	51.2	52.6	47.2	53.7	16.5	48.0	49.6	45.7	43,1	40 40 50	15.0	42.2 6	40.6	40.6
	-	-		dighest during the w	-	70.7 5	72.0 5	79.2 4	55.1	68.9	68.0	67.5	72.5	71.7 4	73.1 4	69.2	69.5	65.1 4	85.1
	-0"	3	renhei	72 observations, co	inches	650	29.750	29.778	29,902	29.593	29.655	29.512	29,675	30.022	876	198	803	\$9,935	29.762
	mor	1 ,191 bus	Sarome Patogra	Mean height of the localions, co	ine	. 29	29	29	23	28	53	20	29	90	29	29.	<b>8</b>	6	<u> </u>
				01		full '	1 gr.		t gr.	29th, full	d qr.	¥	dr.	-	d qr.	TOT.	qr.		
			Phase	of the Moon	- 1	ne 30tlı,	7th, 3rd	h, new	2211d, 1st	ly 29th	6th, 3rd	14th, nèw	21st, 1st qr. Perigee	28th, full	4th, 3rd qr. Apogee	12th, new in Equator	19th, 1st qr.	h fult	
					_'_	June		15th,		July			-					26th	200
			*	** **		6th	13th	20th	27th	3rd	10th	17th	24th	Sist	ber 7th	14th	21st	28th	lighest west, o weeks.
			184	Week		July	2	2	2	August	20	:	8	â	September 7th	2	2	6	Mean, Highest, or Lowest, of the 13 weeks.
			_			-7				_					02				Z

The Ages of 9 persons were not specified in the Table.
 Inquests are held on violent and sudden deaths which are returned irregularly.

### DEDUCED FROM OBSERVATIONS,

Made at Helston, Cornwall, in Lat. 50° 7' N., Long. 5° 18' W., by M. P. Moyle, Esq.

1844	Mean height of Barometer, corrected and reduced to 32°	of Barometer, corrected and reduced	of Barometer, corrected and reduced	of Barometer, corrected and reduced	of Barometer, corrected and reduced	of Barometer, corrected and reduced		T		OMETER gistering	RS .	Wet bulb			Average Cloudi-		in :
Months		Observations	Highest	Lowest	Difference	Mean of Daily Observations	Mean of 3 Daily Observations	Average force	General Direction	ness.	Days	Inches					
July	30.098	62.7	78	48	30	61.5	58.4	1.03	N.	3.86	7	1.22					
August .	30.005	59.5	71	45	26	58.5	54.6	1.30	N	3.80	10	2.87					
Sept	30.105	60.4	75	42	33	58.5	56.5	1.1	. <b>E</b>	3.63	9	1.34					

Barometer corrected for capacity of Cistern, Capillary Action and Temperature, and not for elevation of 106 feet above sea level.

Observations made thrice a day, 9 A.M., 3 P.M., and 9 P.M.

# DEDUCED FROM OBSERVATIONS, Made at Falmouth, by Lovell Squire, Esq.

YEAR 1844	Mean beight of Barometer	·	Т	HERM	OMETE	W	ind	$\Lambda$ verage	Wet	Rain			
	corrected and reduced to 32°	Mean of 3	Self Registering Wet				Wet bulb	1	Cloudi-		in [		
Months.				Daily Observations	Highest	Lowest	Difference	Mean of Daily Observations	Mean of 3 Daily Observations	force	Direction*		Days
July	29.93	62.6	71	51	20	61	• •	••	N.W. & S.W.	6.2	14	1.400	
August .	29.83	58.59	67	50	17	57.83	55.51	• •	N.W.& W.	6.7	16	2.916	
Sept	29.96	59.11	69	47	22	58.28	55.37	• •	N.E. & N.	5.6	12	1.231	

<sup>\*</sup> From one observation daily.

### DEDUCED FROM OBSERVATIONS,

Made at Black Rock, Brighton, by J. O. N. RUTTER, Esq., F. R. A. S.

YEAR 1844	Mean height of		Т	HERM	Wind		Average	Wet	Rain			
	Barometer corrected and reduced	* Mean of 3	Self Registering In Shade, North Wet				Wet bulb		Cloudi-		in	
Months	to 32° Inches	Daily Observations	Highest by Day	Lowest at Night	Difference		Mean of 3 Daily Observations	force	Direction		Days	Inche
July	29.917	64.7	69.5	54.1	15.4				e •	• •	8	
August .	29.834	61.0	64.7	53.3	11.4	* •	* *	• •	• •	••	10	
Sept	29.994	60.8	64.4	53.5	10.9	••	. 2.	• •	• •	• •	8	••

<sup>\*</sup> Observations made 3 times daily, 9 A. M., 3 and 9 P. M.

# DEDUCED FROM OBSERVATIONS, Made at Truro, Cornwall, by Dr. C. BARHAM.

YEAR 1844	Mean height of Barometer.		T	THERM	Wind			Wet	Rain			
	eorrected and reduced			Self Registering Wet bulk						Average Cloudi-	Days more or	in
Months.	to 32° * Inches		Highest	Lowest	Difference		Mean of Daily Observations	force †	General Direction	ness.	less	Inche
July	30.093	Absolute Mean	77 69.6	42 51.5	35 18.1	60.5	:	1.21	W.N.W	4.2	17	4.17
August .	30.008	Absolute Mean	68 64.2	39 47.8	29 16.4	55.5		1.40	W.	5.4	18	5.18
Sept	30.139	Absolute Mean	73 67.7	35 48.8	38	58.2		0.87	E.]_	4.92	8	3.16

These observations are uncorrected, the mean temperature of Morcury for each month was as follows:—July, 65.3; August, 62.5; September, 65.2; the observations were made at 9 A.M., 3 P.M., and 9 P.M. † Estimated loosely by a scale from 0 to 6, indicating the varying force from calm to violent storm.

### STATE OF THE PUBLIC HEALTH IN THE LAST QUARTER.

"The Quarterly Returns are obtained from 115 Districts, sub-divided into 576 Sub-Districts. Thirty-four Districts are placed under the Metropolis, and the remaining 81 Districts comprize, with some agricultural Districts, the principal towns and cities of England. The population was 6,578,912 in 1841."

The deaths registered in the last quarter, (ending September 30th,) were 38784; or 141 less than the deaths (38925) in the June quarter; and 1701 more than 37083, the average of five previous September quarters. Allowing for the increase of population the mortality will be found to be about the average of the summer quarter (July, August, September); that quarter being now the least fatal in the year.

The deaths in the Metropolis amounted to 11825; which are 354 more than were registered in the Spring quarter, and 694 above the average uncorrected for the increase of population. Small Pox and Scarlatina were both epidemic. The deaths from Small Pox in the first week of the quarter were 36, in the last 37; from Scarlatina 58 in the first week, 85 in the last, and 99 in the tenth. Measles was fatal but to a small extent; Typhus shewed a tendency to increase for the deaths from this disease, including what is sometimes called "common continued fever," and "fever," amounted to 424; while the average is 352.

In the country districts the most remarkable increase of mortality occurred in Cornwall; in the Redruth and Penzance districts the following were the results of registration:—

	Annual	Deaths
	Average Deaths	in
	in 5 Summer	the last
	quarters.	Quarter.
Redruth	218	. 419
Penzance	216	. 475

Measles and Hooping Cough prevailed in both the districts. Small Pox has been epidemic in many districts all over the kingdom, and has proved fatal to great numbers, whose vaccination had been neglected.

The return includes the deaths of *nine persons* who were suffocated or trampled to death at Nottingham, whilst witnessing a public execution. At Stockport six persons were accidentally killed by the falling of a wall during a heavy storm of rain.

The mean temperature at the Royal Observatory, Greenwich, was 58° 9 degrees, or 2 degrees lower than the mean temperature of Greenwich on an average of twenty-five years. The mean of the highest temperature of day was 66° 9, the mean of the lowest temperature of night 51° 4; the difference 15° 5. The highest temperature attained in the sun was 115° in the week ending July 27th; the lowest temperature on the grass in the same week was 40° 5; the temperature of the air ranged in the same week from 85° 1 to 53° 7. In the 13 weeks the lowest temperature of the air was 40° 6; of the grass on the ground 30° 0. The difference between the dew point and air temperature was 5° 3. The fall of rain was 5° 38 inches.

Districts in which the mortality was greater than the mean mortality of the corresponding quarter in the same district. The west, north, and south district of the Metropolis, Portsea Island, St. Albans, Wycombe, Bedford, Colchester, Plymouth, Redruth, Penzance, Stroud, Hereford, Shrewsbury, Kidderminster, Wolverhampton, Birmingham, Aston, Coventry, Nottingham, Liverpool, West Derby, Blackburn, Chorlton, Bradford, Merthyr Tydfil.

Districts in which the mortality was less than the average mortality of the corresponding quarter in the same districts. Central districts of the Metropolis, *Norwich*, Devizes, Exeter, Basford, Stockport, Macclesfield, Great Boughton, Bury, Prescot, Salford, Ashton, Sheffield, Hull, *Sunderland*, Tynemouth, Newcastle-on-Tyne, Pontypool, Wrexham.

### REGISTRARS' NOTES.

The following Direction was addressed to the 447 Registrars in the Country:-

"If at any time the number of deaths registered during the Quarter has been above the average, state, in a note at the foot of the Return, whether any epidemic disease, such as Measles, Typhus, &c., has been prevailing in the District; or if there be any other known circumstance which will account for the increase."

In compliance with the above direction, 178 Registrars have appended explanatory notes to their Returns; the remaining 269 Registrars apparently observed a decrease, or no obvious increase, in the mortality.

An Abstract of the Notes to the Returns is subjoined.

SUP. REG. DISTRICT.	REG. DISTRICT.	
MAIDSTONE		This return is considerably below the average; my district has been healthy during the last quarter.
BRIGHTON	The Palace	Only 2 more deaths this quarter, than were registered in the corresponding quarter last year; my district continues very healthy.
		The smallest number registered since the passing of the Registration Act; and 21 less than the corresponding quarter last year.
PORTSEA ISLAND	Kingston	In my last return I informed you that there was an increase in the epidemic diseases, (compared with corresponding quarters) in my district; and they have continued to increase during this quarter also, there being 19 more deaths than in the corresponding quarter of 1843. Measles continues to rage, 11 cases having proved stal; Small Rox has been very prevalent, 4 cases of death having been registered, and all without previous vaccination; Scarlatina has also been very general, but only 2 cases have proved stal. I have also registered I case of English Cholcra.
		Six of the deaths are from Scarlatina; otherwise, the average of deaths would be much smaller than usual.
WINDSOR	Egham	About the average; district generally healthy.
ST. ALBANS		The number of deaths registered this quarter is 19 above the corresponding quarter of last year, occasioned chiefly by inflammatory diseases among children.
Namedon institution distributions included a	Harpenden	The deaths during this quarter shew an increase of about 25 per cent. as compared with the corresponding quarter of 1843. <i>Measles</i> and <i>Typhus</i> have been fatal in my district, which probably accounts for the increase.
WYCOMBE	High Wycombe	The number of deaths this quarter is 24 above the corresponding quarter of 1843, principally owing to <i>Consumption</i> and deaths among infants, as I am not aware of any epidemic in my district.
		The mortality this quarter is less by 15 than in the preceding one, and 2 less than in the corresponding quarter; 1 death of a child was from Small Pox, (natural); district very healthy.
NORTHAMPTON	Bugbrook	The number of deaths this quarter is above the average, Scarlatina prevailing.
		The number of deaths is just half of what it was in the last quarter, ending 30th Junes.
- consumption references estimated estimated entitles of a	St. Giles	Deaths 87.—This district has been very healthy during the quarter; 9 of the deaths were Coroner's Inquests.
BEDFORD	Bedford & Cardington	Number of deaths below the average during the last quarter.
	Bedfard & Kempston	The number of deaths registered this quarter is about one third greater than the average registered in the corresponding quarters of the preceding seven years. Of these deaths, about one-third has been attributed to Diarrhaa and Hooping Cough, which have been very prevalent, another third to Consumption, Atrophy, and Marasnus, and of the remaining third, about one-third part to untimely birth.
CAMBRIDGE	St. Andrew the Less	The number of deaths this quarter is 60, which is about the average; there have been only 4 fatal cases of Small Pox, (all without previous vaccination). The district may be considered healthy.
NORWICH	East Wymer	Small Pox has again appeared in this district, and has been fatal in 9 cases.
		The number of deaths registered during last quarter has not been above the average; 10 deaths from Phthisis, and 6 from Typhus; but no epidemic disease has been prevailing in the district.
	Costany	· ·
		This return is below the average; the district has been remarkably healthy.
		The above number is considerably below the average, the district being very healthy.
EXETER	St. Sidwell	Fifteen less than in the corresponding quarter of last year.
		Five cases of Typhus, (1 of them preceded by Ervsipelas), 2 other cases of Hever 1 Debility following Fever, 2 of Erysipelas, and 2 Inquests.
1		An increase of deaths this quarter above the corresponding quarter of last year; 4 cases of Coroner's Inqests, 13 children died from Diarrhœa, Thrush, and Hooping Cough.
	East Budleigh	Erysipelas has prevailed and caused 2 deaths; 5 deaths also from Phthisis.
3.5 (1) (1.5 (1.5 (1.5 (1.5 (1.5 (1.5 (1.5 (1.5	Heavitree	The number of deaths this quarter is much the same as the last; Natural Decay 4, Disease of the Lungs 8, Insanity 1, Hooping Cough 2, Disease of the Womb 2, Fever 3, Disease of the Heart 1, Diabetes 1, Jaundice 2, Paralysis 1, Enteritis 1, Rupture of Blood Vessel 1, Inquest 1,—28.
	Alphington	My district is particularly healthy, shewing the smallest amount of deaths since the Registration Act.

SUP. REG. DISTRICT.	REG.DISTRICT.	The last was last the same and
ST. THOMAS		The deaths are about the average; district very healthy.  The mortality this quarter is 47 above the average, which is partly accounted for by the prevalence of Mcasles and Diarrhea,—19 deaths from the former, and 11 from the latter; the remaining 17 are from various causes.
	Charles the Martyr	The mortality is above the average, which may be attributed to the prevalence of Measles and Small Pox: the cases of the latter epidemic were all unvaccinated.
		Measles and Hooping Cough have been very prevalent
	Redruth	The deaths in this quarter exceed those in the corresponding quarter of 1843 by 57, arising chiefly from the prevalence of Measles.
	Illogan	The number of deaths this quarter is nearly double the amount of the average of the four preceding quarters; which increase is caused by the <i>Measles</i> among children. Out of 80 deaths there were 51 children under five years of age.
	Camborne	The amazing increase of deaths arises from the prevalence of Measles and Hooping Cough; above 50 have occurred from these diseases and their consequences. An unsual number of cases of Phthisis also appears, amounting to 17. The average of deaths taken above a year since was 49.
		About 20 above the average, chiefly owing to the prevalence of Measles in my district.
PENZANCE	Penzance	The number of deaths this quarter has been considerably above the average, owing to the great mortality by Measles and Hooping Cough.
	Saint Just	The number of deaths this quarter, being 49, (omitting 14 Coroner's Inquests, some of which happened in the preceding quarters,) is above the average; which is apparently caused by <i>Measles</i> among children, as 26 have happened from the effects of that epidemic.
	St. Buryan	This quarter is above the average of preceding quarters by 7.—Deaths 20: Debility 5, Hooping Cough 4, Phthisis 3, Inflammation on the Chest 2, Stoppage in the Stomach 1, Convulsion Fits 1, Measles 2, found downed 1, Mortification 1; district healthy at present.
<del>edicides, aggressistation</del> — 4 to 70 to 7	St. Ives	Much above the average, owing to the prevalence of Hooping Cough and Measles concurrent. 51 of the number are children who have died of those diseases.
	Uny Lelant	Measles and Hooping Cough have been very prevalent in my district the last 2 months of the past quarter. It shows an increase of 15 above any other quarter since the commencement of the Registration Act.
andresting controller in the controller of the self of	Marazion	The number of deaths registered this quarter is 28, a few deaths above the average, occasioned by Measles and Hooping Cough prevailing.
BATH	Lansdown	A little above the average.—Hooping Cough 8, Scarlet Fever 4, Measles 2, Small Pox, without previous vaccination, I.
0 0 0 0 0 0 0 0 0 0 0 0 0	Batheaston	The number of Births and Deaths registered by me during the last quarter is much below the average; the registered causes of death are as follows:—Consumption 9, Dropsy 3, Decay of Nature 3, Epilepsy 3, Paralysis 2, Debility 1, Disease of the Liver 1, Water on the Brain 1, Teething 1, Measles 1, Searlet Fever 1, Disease of Intestines 1.
BRISTOL		One death from Small Pox without previous vaccination.
# # # # # # # # # # # # # # # # # # #	St. Augustine	About the average for the quarter, but the deaths during the month of September have been more than double the number of those registered in either July or August, being respectively 18, 17, and 37, though there does not appear to be any particular disease prevalent.
6 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	St. Mary Redcliff	The number of deaths is not above the average; there have been 15 deaths from Small Pox, only one of which had been previously vaccinated.
g o p. a la talla .a . a . e . a .	St. Paul	During this quarter 5 deaths have been registered from Hooping Cough, and 7 from Small Pox; all without previous vaccination.
CLIFTON	Ashley	The number of deaths registered this quarter is 12 above the corresponding quarter, and II under the previous quarter; epidemics have been prevalent; 6 deaths have been registered from Scarletina, 3 from Hooping Cough, 5 from Measles, and 1 from Typhus.
gantificacional national in garage (a a a a a a a a a a a a a a a a a a a	St. George	The number of deaths this quarter (including 4 inquests not yet registered) is 25 per cent, above the general quarterly average, and 39 per cent, above the corresponding quarter of last year.—Fever, Diarrhea, Hooping Cough, and Measles have prevailed, and the increase is caused by these diseases.
		A few cases of severe Diarrhoa supposed to be caused by fruit.  Decline and Consumption 8, Dropsy 2, Fits 2, Scarlatina and Typhus 7, Old Age 2, Inquests 4.
		The number of deaths is 10 more than in the corresponding quarter for 1843; no epidemic disease has been prevailing in the district.
SHREWSBURY	. St. Chad	Small Pox still continues in this district and town, although all means in their power has been taken by the different parishes to arrest its progress.
	St. Mary	I have this quarter registered 110 deaths, the largest number registered by me, with one exception, since the operation of the Act; out of which number are 26 cases of Small Pox—25 "without previous vaccination" and 1 only "with previous vaccination;" quarterly average about 83, the Small Pox cases accounting for the increase. Some stringent measures should be adopted to compel parents and guardians of children to take them for vaccination; their negligence is criminal. The largest parish in my district has adopted the Provisions of the Vaccination Act, 3 & 4 Vic., c. 29; yet few (taking into consideration the population) have availed themselves of the benefits thereby offered. Other cases of epidemics are, Measles 7, Typhus 1.
WORCESTER	North	The number of deaths in my district is rather more than the usual average; 4 deaths from Typhus have taken place, all of which occurred in the Worcester Infirmary.
KIDDERMINSTER	Kidderminster	The cases of Fever and Consumption appear to be very considerably beyond the usual number.
DUDLEY	Sedgley	The number of deaths registered in my district this quarter is 121, being 30 less than in the preceding one; out of which number 25 (chiefly children at various ages under 7 years) have died from Diarrhæa, or Bowel Complaint, and Diarrhæa Convulsions.—From Consumption 10, Typhus 6, and Measles 1.
eurge for intoning a spire in	. Dudley	Deaths registered in my district this quarter 179, being 11 more than the last: Bowel Complaint has been rather prevalent. There have been 29 deaths from bowel complaint,

SUP REG. DISTRICT. REG. DI	STRICT.
WOLVERHAMPTON Eastern	The number of deaths registered by me this quarter is greater than the corresponding
	ones of 1842 and 1843, showing an increase of 12 over the former, and of 30 over the latter, and an increase of 30 over the average, owing to the prevalence of Diurrhaa and Hooping Cough among children.
Western	I have an increase of deaths this quarter; 14 more than the corresponding quarter of 1843, and 33 more than last Midsummer quarter. I have registered this quarter no less than 23 cases of Bowel Complaint, which no doubt is the cause of the increase.
Bilston	The increase in the number of deaths this quarter is owing to the prevalence of Diarrhæa and Inflammation of the Bowels among young persons.—There has been no epidemic.
Willenhall.	The number of deaths this quarter is 4 more than in the corresponding quarter of 1843.  The increase is caused by Inflammation in the Bowels and Consumption.
WOLSTANTON Burslem	
BIRMINGHAM St. George	
St. Paul	The number of deaths this quarter is 14 above the yearly average and 35 above the corresponding quarter of 1843. There is no epidemic in this district, but the mortality has been principally among children under the age of twelve months, making rather more than one third of the total number. Much of this, I fear, is attributable to want of proper care and nourishment among the children of the poor, and sometimes to debility of parents caused by the employment and confinement of mothers in manufac-
St. Philip	tories, and consequent inattention to cleanliness, and premature labours.  The deaths this quarter are 22 more than the corresponding quarter of 1843, and 14 more than the last quarter.—15 deaths by Consumption, 4 by Scarlatina, and 3 of
Lady Wood	Small Pox (1 of which was a second attack) without previous vaccination.  The number of deaths this quarter presents an increase of about 10 per cent, above the preceding one, but the number of epidemics a decrease from 22 to 9. A continual increase of population in the district may have produced the former.
ASTON Sutton Coldy	field The increase arises from having registered eleven deaths upon which inquests had been held previous to the commencement of this quarter's return.
COVENTRY Holy Trinit	The dispressions to the commencement of this quarter's return.  Of the above return of 82 deaths in my district 44 are children 1 year old and under, of whom 15 died of Disease of the Bowels or Chest, 11 of Natural Weakness, and 5 of Convulsions.
LEICESTER East	******** There is an increase of 17 deaths over the preceding quarter, though I consider it about an average number. Scarlatina is very prevalent at this time; I have registered 20 deaths from that epidemic during the quarter; in other respects I consider the district
West	healthy.  The district during this quarter very healthy, not any epidemic prevailing; the number
LINCOLN Home	registered this quarter is less by 10 than the preceding quarter.
	having been prevalent except Consumption which has done its work, numbering one fourth, viz., 22, as it victims. I repeat my opinion that to our want of drainage much of Consumption is to be attributed.
	Although the number of deaths in my district during the past quarter is less by 4 than in the preceding one, yet I must again notice the prevalence of Typhus the number of deaths from it amounting to one sixth of the whole, the cases having been more severe and the disease most fatal in those houses near which the sewers are left uncovered, and from which the drainage is bad. This has been the case in the village of Normanby, especially where the principal drain having no outfall, accumulations have taken place in the smaller ones producing Malaria which gave rise to the disease in the first instance, and has been the means of its continuance.
NOTTINGHAM St. Ann	There are 26 deaths above the average this quarter; 19 cases of Phthisis have been registered; also a few cases of Small Pox. Although three Surgeons have been appointed as Vaccinators, totally distinct from the Medical Officers to the Union, yet a great majority of the working class are so utterly regardless of their childrens' welfare as not only to reject the proffered boon, but frequently to insult the surgeon appointed to carry the law into effect, although the children are vaccinated gratuitously.  In 1837 From 1st July to Sept. 30 the deaths were 70  1838
Exchange	This quarter's return is more than an average, owing to 9 persons having been suffocated or trampled to death whilst mitraging a with a wife or trampled to death whilst mitraging a with a wife or trampled to death whilst mitraging a with a wife or trampled to death whilst mitraging a with a wife or trampled to death whilst mitraging a with a wife or trampled to death whilst mitraging a wind or transport with the wind of the wind or transport will be with the wind of the wind or transport will be will be wind or transport will be will be wind or transport will be will
BASFORD Bulwell	The return shows an unusually small number of deaths
DERBY St. Alkmund	The number of deaths this quarter is 16 above the average, and also 6 above the corresponding quarter of last year; but I am not aware that any particular epidemic has prevailed.
STOCKPORT Stockport (S	Second) The deaths registered this quarter are less by 30 than the lowest average. My district appears to be in a very healthy state.
Hyde	•••••• Deaths are below general average.
•••••• Heaton Norr	Below the average; and although 20 less than its correspondent quarter, 1843, it shows an increase of 7 per cent. over it, in 9 principal diseases. The increase is 8 per cent. in Diarrhoea, 5 per cent. in Convulsions and Debility from Birth, and in Dysentery about 1 per cent.; whilst there is a decrease of 5 per cent. in Phthisis and Scarlatine, nearly 2 per cent. in Pneumonia, more than 2 per cent. in Premature Birth, and a total absence of Fever.
MACGIESERIA D	There are 14 more deaths this quarter than the last, and more than the corresponding quarter of 1843; 8 under 12 months; 6 were accidentally killed by the falling of a wall during a heavy storm of rain. The district at present is healthy.
MACCLESFIELD West	The average number of deaths in the quarters ending September, for the seven years during which the Act has been in operation, is 72·42; the number for the last quarter is 73.  Inflammation of the Chest and Bowels, Diarrhæa, and English Cholera have been prevalent, of which a few cases have terminated fatally.

SUP. REG. DISTRICT	REG. DISTRICT.	
	Sutton	The number of deaths in the present quarter is very much below the average.
	Bollington	The number of deaths registered during this quarter is very considerably below the average, being only 33 in number, 11 of which were children under 1 year.
		Six of the deaths that I have registered this quarter, have been from Phthisis.
LIVERPOOL		The number of deaths in this return is above the average; Consumption, Diarrhoea, Infammation of the Lungs and Bowels, having been the prevailing diseases.
	Great Howard Street	This return shews an increase on the corresponding quarter of last year of 88. The whole of my district is at present in an unhealthy state. Small Pox has decreased though it has not entirely disappeared; the weather has been most variable, and unfavorable to invalids and children.
	Dale Street	An increase of 56 deaths more than the previous quarter has occurred in my district, principally amongst children afflicted with Convulsions. Teething, Measles, and Inflammation on the Lungs.
1210000	St. Thomas	The deaths this quarter are 2 more than the corresponding quarter of 1843. The principal diseases are Phthisis 12, Enteritis, &c. 45, Fever 20, Premature Births 10, Hydrocephalus 12, Teething 13, Pneumonia 22, Measles 10, Croup 4, Delirium Tremens 4, Inquests 11.
6,0000	Mount Pleasant	The number of deaths in my district this quarter is 267, the smallest number I have ever registered during the $3\frac{1}{4}$ years I have been Registrar; in comparing this quarter's deaths with the corresponding quarters of the 3 last years I find they are as follows, viz.:—September quarter, 1841, deaths 298, September quarter, 1842, 329, September quarter, 1843, 290, so that I must conclude that my district is in a very healthy state, and has been so for the last 6 months.
000000	Islington	There is an increase of 20 deaths in this quarter as compared with the corresponding quarter of 1843, principally caused by <i>Enteritis</i> ; I have registered 13 from that disease. I consider that the district continues healthy.
	Saint George	The deaths this quarter are below the average, although Small Pox has prevailed very much in the district, and has been fatal in 12 cases.
WEST DERBY	West Derby	The mortality this quarter does not exceed the general average, although Pulmonary Disorders, Scarlatina and Typhus, Small Pox, and Dysentery, have been very prevalent. Scarlatina proved fatal in 10 cases.
	Crosby	A decrease of 6 deaths compared with June quarter, though Typhus and Scarlatina have been prevailing diseases in my district.
468000	Toxteth Park	There is an increase of deaths this quarter (58) compared with the former quarter; Typhus, Scarlatina, and Measles are prevalent in my district. Typhus 6, Scarlatina 10, and Measles 9.
BLACKBURN	Blackburn	The number of deaths this quarter is about the average; no epidemic prevailing except Small Pox: 37 children have died of that complaint, 27 without previous vaccination, and the remaining ten of a second attack after vaccination; 66 persons have died of Consumption, and 43 (infants) of Fits.
.,,,	Mellor	The number of deaths this quarter is 19 less than the former quarter, and below the usual average.
00000	Darwen	We have more than an average number of deaths in this quarter; the prevalence of <i>Hooping Cough</i> accounts for the difference.
	Great Harwood	This quarter is above the average.
	Oswaldtwistle	The return is 16 less than the corresponding quarter, 1843, and 9 less than the previous quarter.
J# 6 6 6 8	Witton	I have a decrease of 5 deaths in my district this quarter as compared with the previous quarter, although Measles has been and is prevailing.  There are 20 deaths less than the coverence dier quarter less trees. I have hed 20 deaths
PRESTON		There are 20 deaths less than the corresponding quarter last year. I have had 20 deaths from Small Pox without previous vaccination.
	Longton	The number of deaths registered by me this quarter is 24, being 7 less than in the corresponding quarter of last year, 9½ less than an average of the four preceding quarters, and 16 less than an average of the last 3 years, which is as far back as I am able to refer. In the village of Longton, Small Pox is very prevalent; hitherto, however, no cases have proved fatal. With the above exception my district is in a healthy state. Of the 24 deaths, 5 are of children under the age of 1 year, and 6 of persons who have respectively attained the ages of 67, 73, 77, 77, 77, and 84 years. The causes of death are Phthisis 8, Convulsions 3, Apoplexy 2, Small Pox 1, Worm Fever 1, Cirrhosis 1, Croup 1, Pneumonia 2, Thrush 1, Debility from Birth 1, Disease of the Heart 1, Old Age 2.
9999999	Walton-le-dale	The quarter's return (36) is 3 above the corresponding quarter of 1843, and one-eighth under the average for the September quarters since the commencement.
	Spotland	Sixteen less than last quarter; consequently my district is remarkably healthy.
BURY	Bury, South	There has been about one-seventh fewer deaths in my district during the past quarter than in the corresponding quarters in the last 6 years, although 9 of the 55 deaths have been from accident, and there has been a very considerable increase of population since 1841.
••••••	Birtle	This district is healthy
		The number of deaths is 8 below the average.
	Holcombe	The number of deaths, viz. 22, is exactly the average of the corresponding quarters since the commencement of the Act.
POLZOP		Below the average for the corresponding quarters.
BOLTON		There was an increase of deaths during the last quarter, owing to Convulsions among children.
	Little Bolton	The number of deaths this quarter is below the average of corresponding quarters. This district is in a very healthy state; there being no particular disorder prevalent, and I believe that it is chiefly owing to the improved condition of the working classes. I have had I death from Small Pox, not vaccinated, and 2 of Typhus during the quarter.
***********	Horwich	The number of deaths in this quarter is below the average; the district is in a very healthy state.
**************************************	Tong with Haulgh	The number stated is considerably below my usual average, and in that small account are included 4 persons, whose several ages are 66, 71, 80, and 100 years.

SUP. REG. DISTRICT	REG. DISTRICT.	
WIGAN	Wigan	Small Pox, which has been so prevalent here since last Autumn, has now nearly disappeared, only 15 cases have been registered during the past quarter.
B	Standish	It is a remarkable coincidence, that the number of deaths is the same as that of 1843 for the same quarter; the numbers, however, are reversed, the males being 14, the females 12.
PRESCOT	Aspull	An average decrease of deaths this quarter of 8, compared with the preceding quarters. This district has been very healthy this quarter, 10 below the average of the corresponding
	Prescot	quarters of 1841 2-3.  This district has been remarkably healthy during the past quarter, the number of death being under the general average; and though we have had a great many Irish labourer lodging in the town during the harvest, still the mortality has not been increased by that circumstance, there being only two more deaths this quarter than the preceding one.
		The number of deaths is much below the average.  The number of deaths registered during the last 3 months has been about 10 less than the average quarter, and about 30 less than the average summer quarter, and 25 greate than the last quarter, which was an unusually healthy one. Trade has been verbrisk, provisions cheap, and the weather fine, for the most part dry and warm, wit constrable movement of the air. The mortality would have been low but for the pre valence among the children of the poor of Diarrhea, and other intestinal disease. 3 of the 173 deaths registered have been those of young children, caused by such disease. 20 of which occurred in houses of the lowest class, and 12 in streets of the lowest class and worst condition. About one-third of the inhabitants of this district reside in house of the third class; 20 of the 33 fatal cases of Diarrhea, &c. have occurred among them. About one-fifth inhabit streets of the worst class, among whom 12 out of 35 of these cases have happened.
6 0 0 0 0 0 k.c.	Ardwick	The number of deaths in this district is greater than usual, occasioned principally be Consumption and general Chronic diseases. There are as many as 147 deaths this summer quarter, and in the corresponding one of 1842, there were only 85.
\$ 6 6 6 6 6 C.S.	Barton upon Irwell	The deaths in my district are about an average of last quarter; I have had 2 fatal cases of Typhus and 2 of Cholera.
MANCHESTER	Ancoats	Total deaths 290.—Scarlatina 4, Measles 6, Inflammation of Lungs 17, Inflammation o Bowels 5, Consumption 25, Fever I, Croup 4, Childbed 4, Dropsy 7, Convulsion within first month 16, Convulsions within 3 months 25, Convulsions within the year o upwards 16, Teething 19, Small Pox without previous vaccination 2, Coroner's cases 12 General causes 127.
	St. George	The number of deaths registered in this district during the present quarter is below the average. The principal causes have been,—Convulsions 33, Consumption 27, Dentition 21, Pneumonitis 15, Measles 14, Debility from Birth 14, Diarrhoa 11, Tabes Mesen terica 9, Hydrocephalus 9, Childbirth 7, Disease of the Liver 6, Small Pox, nature 2, English Cholera 2.
. Энефиянциянняя разричинамического в в в в	Market Street	Independently of the Workhouse, Infirmary, and Coroner's Inquests, the deaths in my district in the quarter ending 30th September last are 136, being 41 less than in the corresponding quarter of 1843.
	London Road	The total number of deaths is 246—Males 127, Females 119. The number of death registered, exceeds that of the last (June) quarter by 52, and is less than that of the corresponding quarter for 1843 by 53. The most prevalent diseases have been Diarrhead 40 cases, as in the preceding September, and Scarlatina. Measles and Hooping Coughave been less frequent. Fewer cases of Fever by one-third; deaths arising from Infantile Debility and Convulsions are fewer, as 52 to 21.
		The above is about 50 less than the average of the corresponding quarters. Number of deaths in July 71,—August 63,—September 73.
ASHTON	Ashton Town	Above the average of the corresponding quarters. I have not before noticed the death of females to exceed the males as they do in this and the quarter preceding, via about one-fourth in each. 3 males and 4 females died from Small Pox without previous vaccination, and 1 female 7 years old after having been vaccinated six year ago.
	Audenshaw	The number of deaths this quarter is 3 less than the corresponding quarter of 1843. But there is an increase of deaths arising from Premature Birth.
		The number of deaths registered by me, is about an average of the last two corresponding quarters. There is no prevailing epidemic in this neighbourhood at present Consumption has been fatal in 5 cases, being a fifth of the whole number.
		Much below the usual average.
		The number of deaths is much below the average.  There are 33 deaths more than the corresponding quarter of 1843, and 36 deaths less
		than the September quarter of 1842; there has been no particular Epidemic.
	Middleton	The return is one-fourth below the average.
	Royton	The deaths are 4 and one-third below the average of three corresponding quarters. This includes 3 Inquests, which, if deducted, would bring the mortality from disease still lower. Bowel complaints, and Inflammatory diseases of the Chest, have prevailed but Small Pox has been most prevalent.
SHEFFIELD	South.	The mortality during the quarter is below the average.  The amount of deaths is about an average, though Measles has carried off 9, and Small
	Kirkheaton	Pox (principally without vaccination) 10.
Approximate to the second seco	3.5	The return exhibits the lowest number of deaths that has occurred in this district since the Registration Act has been in operation.
	Honley	Fifteen deaths are considerably less than the smallest number I remember to have registered in any previous quarter for this district; during the corresponding quarter of last year 45, and during the quarter ending June 30, 1844, 38, and for the last quarter ending 30th September, 1844, 15.
with the sharehoused admiration and statement admiration of a	Lockwood	Not quite an average. On looking over the Registry I find that one half of the deaths which have occurred during this quarter, are infants under 20 weeks old. I find that the average age does not exceed 13½ years. Diarrhæa and other affections of the bowels are prevailing to a considerable extent in this district at the present time, owing in a great measure, perhaps, to the very limited supply of good and wholesome water, occasioned by the dryness of the season.

SUP. REG. DISTRICT.	REG. DISTRICT.	The number of deaths is much below the average.
		The above is a little below the average of the corresponding quarters.
	Elland	The number of deaths registered this quarter is one less than the corresponding quarter last year.
900000000000000000000000000000000000000		Rather below the average; district very healthy.  I have registered 26 deaths this quarter, being 2 more than the quarter ending September, 1843.
	Sowerby	My district is very healthy, and the deaths in this quarter are considerably below the average—fewer than at any former period since I commenced registering.
BRADFORD	East-end	The deaths in my district in the quarter ending September, 1843, were—Males 75, Females 55.—Total 160; the deaths in the present quarter ending September, 1844, are—Males 91, Females 72.—Total 163; shewing an increase of only 3 deaths, whilst the population of my district within the last 12 months has rapidly increased.
		The number of deaths is not above the average, though Scarlatina has prevailed much during the month of September, and has proved fatal to a great number of children.
		Scarlatina and Typhus have prevailed in a slight degree, otherwise the district has been peculiarly healthy.
	Horton	Notwithstanding there being no increase of deaths the Small Pox has prevailed in this district as throughout most of England, and 10 fatal cases, 7 without and 3 after vaccination have occurred, besides 7 of Sarlatina, which has also prevailed very generally.
	•	The deaths are 19 above the average of the 7 corresponding quarters; the increase has been from <i>Phthisis</i> and <i>Hydrocephalus</i> , and in cases of inquests. Phthisis has been fatal in 13 cases, being 7 above the average.
		The number of deaths registered this quarter is 30 more than occurred in the corresponding quarter of last year, but not more than the average of former years. Diarrhea and Hydrocephalus have been the most prevalent, and principally among children.
•••••••	Whitkirk	Out of the 19 deaths 10 were infants not exceeding 6 months old.
		Ten cases of Small Pox natural, 1 case of Small Pox 3 years after Cow Pox, 2 cases Scarlatina, 1 Measles.
		Considerably below the average.  The number of deaths registered this quarter is 14 less than last, but 22 more than
000000000000000000000000000000000000000	rr aimyute	the corresponding quarter of 1843. Small Pox and Scarlet Fever have been prevalent during last quarter and part of this, principally among children; there have also been a few cases of Croup and Typhus. The deaths from epidemics are—Small Pox 5, Scarlet Fever 4, Typhus 3, Croup 4, Diarrhea 4—Total 20.
		Below the average.
***************************************		Above the average for the summer quarter, but from no particular cause, the district being tolerably healthy during the quarter.
TYNEMOUTH		My district has been unusually healthy during the quarter, the mortality having been less than at any former period since the commencement of the Registration Act. The corresponding quarters of the six preceding years average 78; the present return of 57 is, therefore, 21 below that average.
		The number of deaths registered this quarter is the usual average. 45 of the number are children under 10 years of age.
	or 6.	This district has been remarkably healthy this quarter, and the number of deaths does not exceed the average amount.
NEWCASTLE UPON TYNE	JAN Sames	My district is at present particularly healthy; it is at least 35 below the average.
LARLISLE		The number of deaths this quarter is below the average. The district in general is in a healthy state at present.
		This return is about 20 per cent. below the average quarterly number of deaths for the September quarter. My district has been particularly healthy, no epidemic nor contagious disease apparently existing.
		I have had 12 deaths this quarter, just the same as the corresponding quarter of 1843, and out of the 12 there are 5 which average nearly 79 years, viz., 83, 70, 70, 76, and 94.
		This district is more healthy than it has been for many previous months, and there has consequently been a decrease in the number of deaths.
and a distribution of the second seco		Rather above the average. Measles has prevailed in part of my district, but only 8 have died of the complaint. I have 29 under 5 years of age, and 13 inquests chiefly from accidents at the collieries.
KENDAL		My district is in a very healthy state at present; out of the 22 deaths 4 occurred at the following ages:—77, 77, 79, and 87 years.
		This quarter is an average of the quarters since the commencement of the Registration, and one-fourth above the average of the corresponding quarters.  Hooping Cough has been prevalent; several cases fatal.
		Rather below the average; and the district is free from any epidemic, and in a very healthy state.
APERCATERING		My district is in a very healthy state at present; there is a decrease of 6 as compared with the corresponding quarter of last year.  My district is the lost by 4. My district.
		'The number of deaths this quarter is not so many as in the last by 4. My district continues in a very healthy state.
MERTHYR TYDFIL		The number of deaths returned is above the average. The increase is attributable principally to the prevalence of Small Pox. Early in the quarter a few cases of deaths from Measles and Scarlet Fever have also contributed to raise the mortality.
		The number of deaths in this quarter is below the average.
HOLYWELL	272.0000 0000000000000000000000000000000	The number of deaths is below the average. Scarlet Fever has prevailed in several parts of the district, and 3 fatal cases have been registered.

# A QUARTERLY TABLE

OF THE

# Mortality

IN A

# HUNDRED AND FIFTEEN OF THE DISTRICTS

OF

# ENGLAND

(INCLUDING THE PRINCIPAL TOWNS,)

SHEWING

THE NUMBER OF DEATHS REGISTERED IN THE SIX YEARS 1838—43, THE AVERAGE

NUMBER OF DEATHS IN THE FIVE AUTUMNS 1838—1842;

AND THE NUMBER OF DEATHS IN THE AUTUMN QUARTER 1844, ENDING DECEMBER 31st.

PUBLISHED BY AUTHORITY OF THE REGISTRAR GENERAL.



LONDON:

HARTNELL, PRINTER RED LION COURT.



### TABLE

OF THE

### MORTALITY

In 115 of the Districts of England (including the principal Towns): shewing the Number of Deaths Registered in the 6 years, the average Number of Deaths in the 5 Autumns, 1838—42, and the Number of Deaths in the Autumn Quarter of 1844, ending Dec. 31st.

		_					-			9 1									1000	101	9
		70	A1. 73		wad in	the we	are	1838-	-42	ing 1844		Fopu-	n	eaths I	Register	ed in th	he years		1838-		ending ending it, 1844
omen I depo	Popu-	Des	iths R	egiste	red in	те уе	HES	Quar • Ave	terly rage	in the Autum rter, ending 31st, 1844	DISTRICTS	lation							Quart • Ave	rage	立 5 mm
ISTRICTS	lation 1841				_	2 1		Five	ive	ns in tarter		1841	œ	6	9	1	2	43	Of Five Years	Of Five Autumns	ths in uarte
		1838	1839	1840	1841	1842	1843	Of F	Of Five Autumns	Deaths i			1.838	1839	1840	1841	1942	1843	53	P S	Deaths ii
			1			1		1			N. Midland								- Lander		
etropolis st Districts	301326	ATTO	6538	6936	6599	6826	7212	1751	1695	1987	division Leicester	50932	1170	1291	1730	1358	1457	1255	350	370	387
	3	8944	7890	8594	8261	8550	9038	2112	2107	2353	Lincoln	36110	620	731	817	800	720	786	184 361	188 386	160 381
	374759		9751	9108	9333	8873	9547	2395	2385	2491	Nottingham	53080	1209	1540	1794	1239 1163	1429	1393 1216	310	316	344
	393247				10004	9947		2640	2683	2999	Basford	59634 35015	1045 769	1200	926	914	909	922	217	219	210
th Districts	479169	13210	11562		11872			2850	2945	3826	Derby	20010	709	002					į	- 1	
Total 1	1915104	53546	46100	47030	46069	46114	49332	11749	11815	13656	N. Western division										
Eastern							į				Stockport	85672	2410	2591	2574	2048	2073	2203	585	548	474
ivision	33210	737	726	700	648	815	774	181	181	204	Macclesfield	56018	1315	1528	1745	1419	1403	1551	370	350	344
ghton	46742	916	898	980	1102	1126	1117	251	266	303	Gt. Broughton,	49085	1040	1237	1384	1246	1189	1091	305	304	263
of Wight	42547	655	630	737	891	721	703	182	175	235	Liverpool	223054	6627	7435	8470	7556	7407	7458	1875	1962	2131
rtsea Island	53036	1264	1180	1140	1211	1152	1342	297	326 110	341 153	West Derby, ?	88652	1625	1746	2206	2202	2405	2312	509	575	814
nchester	23044	421	479	526 402	1	<b>4</b> 69	471 314	116	105	02	(adj. Liverpoot)	75091	1612	2190	2140	1692	1665	2031	465	451	578
ndsor	20502	381	367	-202	550	972	014	31	100		Preston	77189	1726	1812	2637	2028	2050	1938	513	493	429
Midland											Rochdale	60577	1477	1673	1868	1417	1476	1531	396	378	329
Alban	17051	378	310	331	300	340	291	83	84		Bury	77496	1901	1990	2170	2064	1987	1832	506	459	445 621
combe	34150	788		824	751	679	1	187	168		Bolton	97519	2432	2857	2900	2572	260?	2596	668 434	67 l 434	370
ford	19701	359	407		1	414		102		103 181	Wigan	66032	1483	1912	2144	1561	1588	1832 920	260	250	237
rthampton	28103	689	1				1	1 200	161	203	Prescott	43739	916	1056	1155 2399	997	1075 2427	2529	591	574	709
dford	31767	676	1	743 639	\$	604 465		171		132	Chorlton	93736 192408		2418 6774	6489	5821	6151	6263	1597	1612	1653
mbridge	24453	563	547	009	1303	4200	097	144	7.21		Manchester Salford	70228		2449	1998	1971	1977	2098	530	534	525
Eastern											Ashton	173964	1	5489	4873	4282	4786	4391	1196	1112	1173
livision	17790	533	437	488	381	417	393	113	120	122											
chester	25254	694	1	593	499	594	551		1	100	York division								1 000	027	600
orwich	61846	1373	1852	1752		1	t	1 000		9 -	Sheffield	85076	2429		1	1				1	
rmoath	24031	417	440	504	493	516	510	118	119	164	ARTINGCIDACIO	107140		2080			2		1 -00		
Western			1								Halifax	109175		1		3041			1 000		
livision	22130	400	-	48	429	449	633	3 115	114	108	Bradford	132164	1			4360			770	1124	
evizes	23380	1		1	1			.6.2.0			1	41130	1			1206	1042	1			
orchester	31333	1		-	1 000	76	836		1	9.0		47779	1	ļ		1039	1133	1048	274	263	336
. Thomas	47105	1	1				808	21	210	211	Northern							1			
lymouth	36527	86	2 67		1 00	.1		200			division								369	363	270
edrath	48062	91		0.00		100		1 20 21		_	1	56226			-000	1	1	1	1 000		
enzance	4	1	- 1	1 -00	-	1	3				Guitanout	38747 55625	1	968		1		984	0.15	1 000	
ath	69232	156	7 160.	2 130		105	10%	7 76	1	1	Tynemouth Newcastle-on-?		1	į.				1	40	491	37:
Western											Tyne	71850	1	1	á		1	Į.			t .
division	64298	184	2 167	233	0 189	177	1 1749	470	463	3 524	Carlisle	36084	855	739		Ŷ.		§ .	7 7 10 17	min.	
lifton				770	9 151	151	1250		1 -		Cockermouth	_		1	1 000		1	1	304		
troud	00000	95	2 69	3 76	0 72	70		1 200	170	190	Kendal	34694	729	825	821	601	714	666	202		
heltenham	40221	86	9 72		3	1	1	200			Alsway and										
ereford			-	1 0-		ì	1	1			6 A 2	50834	1150	1206	1374	1459	1109	1116	315	274	27:
hrewsbury	21529	1		1 00	l.	3	1	-	1		Donton		1			1	1		147	1 000	
Vorcester				1	1	1					35 57 103	52864		1246	1566	1423	1110	1	1 100		1.0
lidderminster	29408 86028				. 1				1		Mandanna	25958				1	1	1	1 017		
oudley Valsali	1	1	1	1			5				1 197 2		1	ł		1	1		004		
Volverhampton				3	7 215	3 231	9 178	508			Holywell	40787	1				1	575	700		
Volstanton		71							1		1	3810	645	589	632	597	624	0/5	20%		
inumin oh om	13818	335				3			1		0.41 00 . 31	1663308	109321	116505	124664	114664	115834	113869	29049	28867	30302
irmingnam							AL TOP	0370	278	327	I OI CHO METERIODOM	71		1							
ston	5092	· l	00			1		1 -0.	1			-	16.00	20	171004	10.00	18104	115 14363.4	110700	1000	4205
	5092		00	_		1		1 -0.	1			-	162867	16 605	171694	16 733	16194	105201	10798	40682	43958

<sup>\*</sup> The Deaths in the Metropolis for the Years 1840 1-2, have been derived from the Weekly Tables, the Returns for each Year comprising 52 Weeks, or 364 Days. The last quarter in the Metropolis ended Dec. 28, 1844. The Returns from other Places are for the years ending December 31, and the quarter ending Dec. 31st.

<sup>\*</sup> Wandsworth District is included in the return for the Metropolis.

### A TABLE OF THE MORTALITY

IN THE

## Metropolis,

### SHEWING THE NUMBER OF DEATHS FROM ALL CAUSES,

Registered in the 13 Weeks ending Dec. 28th, 1844.

CAUSES OF DEATH	Quarterly Average of Deaths in the 5 Autumns of 1839-43	Deaths in the Autumn Quarter ending Dec. 28th 1844	CAUSES OF DEATH	Quarterly Average of Deaths in the 5 Autumns of 1839-43	Deaths in the Au mn Quarter ending Dec. 28th 1844
ALL CAUSES	12239	13656	IVLaryngitis	.7	17
Specified Causes	12143	13619	Quinsey	1	35
I.—Zymotic (or Epidemic, Ende-)	2355	2991	Bronchitis	178	394
mic, and Contagious) Diseases \$	2000		Pleurisy	22 1317	30
Sporadic Diseases:—			Pneumonia	61	1405 86
II.—Dropsy, Cancer, and other Dis-	7 47 0	1000	Asthma	329	36 <b>6</b>
eases of uncertain or variable Seat	1419	1338	Phthisis or Consumption	1731	1676
III.—Diseases of the Brain, Spinal			Disease of Lungs, &c	187	256
Marrow, Nerves and Senses	1884	2001	V.—Pericarditis	9	35
IVDiseases of the Lungs and of the ?	385 <i>5</i>	4265	Aneurism	9	10
other Organs of Respiration §	2000	4200	Disease of Heart, &c	267	429
V.—Diseases of the Heart and Blood	286	474	VI.—Teething	219	171
VI Discourse of the Stomach Liver			Gastritis '	3	171 15
VI.—Diseases of the Stomach, Liver, and other Organs of Digestion	819	854	Enteritis	208	184
VII.—Diseases of the Kidneys, &c	73	101	Peritonitis	17	29
VIII.—Childbirth, Diseases of the	104	1/10	Tabes Mesenterica	74	101
Uterus, &c	134	17.3	Worms	7	11
IX.—Rheumatism, Diseases of the	69	94	Ascites	12	24
Bones, Joints, &c			Ulceration (of Intestines, &c.)	22	20
X.—Diseases of the Skin, Cellular	18	16	Hernia	24	27
XI.—Old Age	913	898	Colic or Ileus	27 4	29
XII.—Violence, Privation, and In-			Intussusception Stricture	7	8
temperance	318	414	Hæmatemesis	5	9
I.—Small Pox	214	571	Disease of Stomach, &c	55	52
Measles	389 55 <b>2</b>	385 872	Disease of Pancreas	.2	1
Hooping Cough	386	277	Hepatitis	15	23
Croup Thrush	110 59	102 52	Jaundice	27	34
Diarrhœa	122	129	Disease of Liver, &c	95	108
Dysentery	40 .	. 34	Disease of Spleen	1	2
Cholera	7 21	5 32	VII.—Nephritis	7	5
Ague	5	14	Ischuria	3	1
Remittent Fever	5 360	10 385	Diabetes	5	10
Typhus Erysipelas	77	106	Stone	3	9
Syphilis	7	17	Stricture	8	12
Hydrophobia	1	8	Disease of Kidneys, &c	43	60
II.—Inflammation	82 39	40	VIII.—Childbirth	99	121
Dropsy	457	337	Paramenia	3	
Abscess	40 57	61	Ovarium Dropsy	2	9
Purpura	3	6	Disease of Uterus, &c	29	43
Scrofula	26	37 157	IX.—Arthritis	1	
Carcinoma	111	4	Rheumatism	34	43
Gout	12	10	Disease of Joints, &c	34	51
Atrophy Debility	<b>94</b> 263	164 281	X.—Carbuncle	1	1
Malformations	15	17	Phlegmon	1	2
Sudden Deaths	199	186	Ulcer	7	8
III.—Cephalitis	416	160 372	Fistula	. 4	1
Apoplexy	235	311		5	4
Paralysis	213	235	XI.—Old Age	913	898
Convulsions	684 5	699 4	XII.—Intemperance	8	10
Chorea	1	6	Privation	7	10
Epilepsy		54 14	Violent Deaths	303	394
Delirium Tremens Disease of Brain, &c	21 107	25 121	Causes not specified	96	37

# QUARTERLY METEOROLOGICAL TABLE,

November and

1382

30.014 29,738

29,713

44,3

38.7

39.4

32.4

45.1

-9.6

6.0 0.35

365

873

277

385

160

32

1676

25 89

6501

3985

2581

13076

218 1111

18t qr.

11100 23rd 100h 916

Full, 24th

New,

29.695 29.096

45.7

40.8

42.6

49.0 46.0 44.7

0.3 0.3 0.3 0.3 0.1 0.4

4.8 4.3 5.3

+6.4

S.W

7.0

1.58

25 13

23 23

32

499

38

1006

197 480 499 453

163 157 150

949 943

449

877

000

8.2

1.82

70

8.5

7.0 3.0

1.69

36

128

and N.

6.0 4.0

0.5 0.3 0.7

35 36 51

22 2 20 27

88

3 38 27

7th

30.052 30.031

28,0

38.7 32.4

4.3 6.4 6.8 5.0

N.N.E.

830

0.09

9.2

65

84 31

10

160 146 145

255 365 299 224

391 366

278

1170 1326 1150

620 816 478 484

3.0 0.2

6.3 7.2 8.5

31

39 30

26 26

12

815 613

16 40

63

120 105 112 119

203 183 188 195 159 146

286 318 35.8 306 285 305 260 266 286 257

954

257 193 209 146

03

26,0 27.9 33.4 1916 12th

30.8 61.8 65.9 0

加

3rd qr.

0

0

Q

0

0

0

0

46.0

49.0 0

87.2

39.6

5.6 0

11.7

200 200 200 0

+0.3

W.S.W

5.8

0.22

87

16

17

143 132

109 150

508

145

160

[ 879 £66 ; 016

606]

0.08

102

New,

59.1

47.2

54.7 55.7

54.3 55.0

TOTA

Full,

29.672 29,404 29.488 29,611

41.4 42.8 3.0

46.1

44.6

50.I

45.5

29.8

38,1 39.0

3.1

7.8

29.626

40.7

47.6

40.

43.8

41,9

33.0

44.8

ending Week

1844

Mean

Point

Self-Registering

In the Water of the Thames at Greenwich by the Self Registering Thermometer read at 9 o'Clock.

Highest in the sun

Lowest on the grass

MEAN.

THERMOMETERS

Mean height of the Barometer, from 72 observations, corrected and re-duced to 32 deg. Fahrenheit

Highest during the week

Lowest during the week Of the Highest on each day, from 6 observations

Of the Lowest on each day, from 6 observations

Mean of 72 observations

Mean of 72 results

During the week

During the week

Mean of 7 obser-vations

Of the highest on each day from 7 observations.

Of the lowest on each day from 7 observations.

Mean of 72 differences

Mean of the greatest on each day, 6 obser-vations

Greatest the we

Mean, for

The amo

Mean am

Rain in i

Small Po

Measles

Scarlatin

Hooping C

Typhus

Diarrhaa

Influenza

Phthisis

Other dis

0 to 15

15 to 60

60 and up

Deaths fi

Difference

Compiled from the Weekly Tables furnished to the Registrar General by the

,		-	1
the greatest day, 6 obser-	temperature	Differencé be- tween the dew	retur by the
e between the of the week, as ture of the sa age of 25 years	mean t nd the nme we	empe- mean ek on	HOWOLDEE.
irection	Préssure i	W	Er Leog
pressure in k	n lbs. on th	Q N I	
int of Horizoni ir in each week	tal mov	ement	
ount of Cloud,	0-10		
ches (7 days)	· · · · · · · · · · · · · · · · · · ·		
ς			**
aganganyan gabasan dak dalikan bandan dakarah da			the l
and the second s		-	3 wee
ough		,	ks en
outure a l'Attenda de la constantina della const	, .		ling I
, Dysentery, an	d Chol	era.	ec. 28
,			3, 184
r Consumption			, and
eases of the lun	gs		
		Deal Ages, violent	
	eaths)	hs at Thre	,
wards		of len	
om all causes, and sudden dea	(exclus	ive of	
	A Park Base		

Deaths registered in the Metropolis from Small Pox. Measles, Scarlatina, Hooping Cough, Typhus, Diarrhwa, Dysentery, Cholera and Influenza, Consumption, and other diseases of the Lungs; the numbers at each age,\* and the total deaths (except violent and sudden) in each of

# METEOROLOGICAL OBSERVATIONS FOR THE DECEMBER QUARTER, 1844.

	Year	Mean height of			THERM	THERMOMETERS	Ø2		W	WIND			
	1844.	Barometer corrected and reduced to 32°	Mean		Self Re	Self Registering		Wet Bulb	Avelage	General	Average	Wet	Rain
	Months.	Inches	3 Daily Observations	Highest	Lowest	Difference	Mean of daily Mean of daily Observations	Mean of daily Observat.ons	Force		Cloudiness	Days	Inches
Bath C. Brown, Esq.	October November December	* 29·700 29·782 29·970	61 61 61 61 61 61	6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	22 23 23	33 29	61.6 41.7 36.3	<b>4 6 6</b>	8 E E	W.NW.E.NE. E.SE.W.NE.		11 3	2.75 5.11 0.66
Erighton J. O. N. Rutter, Esq.	October November December	29.739. 29.812 30.017.	52.8 45.7 33.46	57.1 49.5 36.2	46.9 42.3 30.6	10.2	52.49 45.78 33.44	* * *	a 'e e	SW. & NE. SW. & NE. NE.	6 6 6	नु हुन	, ,
Carlisle, Harraby Jos, Atkinson, Esq.	October November December	129.600 29.739 30.021	147·1 40·8 31·8	62.5 54.5 43.5	27.0 23.0 14.8	35.5 31.5 28.7	80. 00. 50. 50. 00. 50.	6 6	Moderate 24 days) Strong 7 , , , , , , , , , , , , , , , , , ,	E. 164 days, W. 134 E. 164 days, W. 1134 E. 264, W. 24	v 2 v	138 0.	2.583 1.356 0.234
Derby Mr. J. Davis	October November December	29.55 29.67 29.96	A & A	65 56 42	29 26 18	36 30 24	 6 6 6	3.25 1.70 1.84	96 pt. 65	W. SE. SW. E. E. W. SW. & S. E. & NE.	8 8 8	15 8 4	1.92 3.41 .46
Falmouth LOVELL SQUIRE, Esq.	October November December	\$29.65 29.70 29.87	52.8 48.3 40.7	64 56 51	36 36 35	23 20 20	52:5 47:9 40:5	489 42.9 31.5	\$2.5 1.9 2.4	SW.	57.7 7.7 8:9	20 20 11	4·916 6·179 3·525
<b>Helston</b> M. P. Moyle, Esq.	October November December	29.756 29.758 29.788	53.60 47.76 42.46	67 59 55	35.7	30 24 30	52.44 48.99 43.90	61.28 45.70 40.66	1.33	W. E. & SW. E. & SE.	5.03 5.86 6.23	1 15 16 12	3.70 5.96 2.94
Liverpool Mr. A. Abraham	October November December	29.627 29.759 30.013		61 56 41	30 0 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	29 27 21	49·10 · 43·18 32·17	3.6 1.9	e e e	NW. S. SE. SE.	ဖထက	r~ ∞ €1	2.99 1.27 0.12
Manchester John Leigh, Esq.	October November December	29.56 29.69 29.88		61 55 45	31 288 119	30 27.	48.8 42.3 31.9	a a a	* * * *.	NW. NE.	* * *	2 2 2	2 03 1 :90 0.60
Truro Dr. C. Barham	October November December	29.728 29.798 29.944	6 6	64 49	20 60 F.	3.4 3.4 3.4	49.3		1.2   0.9   1.1	W. by 56, E. by 17, W. by 26, E. by 38, W. by 0, E. by 79,	4200	21 18 11	3.32 7.33 3.76
													The state of the s

\* 2 Observations daily (Sundays excepted). + Observations made at 9 A.M., and 9 P.M. | Estimated approximatively by a gradation from 0 to 6 indicating the Scale from calm to the most violent

### STATE OF THE PUBLIC HEALTH IN THE LAST QUARTER.

"The Quarterly Returns are obtained from 115 Districts, sub-divided into 576 Sub-Districts. Thirty-four Districts are placed under the Metropolis, and the remaining 81 Districts comprize, with some agricultural Districts, the principal towns and cities of England. The population was 6,578,912 in 1841."

The number of deaths registered in the 115 districts for the Autumn Quarter ending December amounted to 43,958; being an increase of 3160 above the yearly average of five years, and 3276 over that of the five Autumns of 1838—42. The total deaths in the year amounted to 167,708; or to 2.4 per cent., making a correction for increase of the Population.

13,656 deaths were registered in the Metropolis for the quarter, and 50,423 for the year, or 2.6 per cent. In the week ending December 21st, 1393 deaths were recorded; being an excess of 403 above the average of the five Autumns, and the greatest number registered in any one week since the Registration Act came into operation. Small Pox and Scarlatina have been epidemic in the Metropolis. Erysipelas and Childbed Fever have also been more than usually fatal. 10 deaths from that rare disease Diabetes were registered. The Registrars in the provinces state, the excess of deaths to have occurred principally from Small Pox, Measles, Scarlatina and Typhus, a great number of deaths have occurred from pulmonary diseases, and been referred to old age. 117 deaths from Small Pox took place at Blackburn; 81 without Vaccination. The Registrars throughout the Country state that the deaths from Small Pox rarely occur after Vaccination performed effectually by qualified Medical Practitioners.

In Dudley 11 persons lost their lives from an explosion in a Coal Pit; at Wrexham 5 persons were killed by an explosion of fire damp. In the district of Oldham 20 persons were killed by the falling of a fire-proof Mill.

Districts in which the mortality was above the average of 5 Autumn Quarters, North, South, East and West districts of the Metropolis, Brighton, Isle of Wight, Norwich, Winchester, Yarmouth, Plymouth, Penzance, Shrewsbury, Liverpool, West Derby, Blackburn, Chorlton, York, and Merthyr Tydfil.

Districts in which the mortality was below the average of 5 Autumn Quarters, Lincoln, Preston, Bolton, Wigan, Sunderland, Gateshead, Newcastle-on-Tyne, and Newtown.

The mean height of the barometer (corrected) was 29.713 inches at Greenwich; the mean temperature of the air 42° 4; the mean dew point 39° 4; the mean temperature of the water of the Thames 44.° 8. The highest temperature of the air was 65° 9, the lowest 21° 6. The thermometer never rose higher than 87° 2 in the sun; nor sunk lower than 7° 9 on the grass. The mean temperature of the air was 2° 6 below the average of 25 years; it was only higher than the average on the 1st, 7th, 8th, week of the quarter; and in the three first weeks of December was so low as 31° 7, 28° 2, 37° 0, being 10° 2, 11° 5, and 3° 4 below the average; the deaths registered in the metropolis were 954, 1150, and 1326, (exclusive of sudden and violent deaths.)

The rain was 9.22 inches; the greater part of which fell in the latter part of October, and in November; the wind in the coldest week was N.N.E., and its average daily horizontal movement during the quarter was 123 miles.

The Meteorological observations from the Country, although not yet taken upon an uniform plan, are in many respects highly interesting; and will, it may be hoped, ultimately end in a connected scheme of systematic observation, with comparable instruments all over the empire. In the mean time Meteorologists are recommended to consult the last Volume of the Greenwich observations, drawn up by the Astronomer Royal.

### REGISTRARS' NOTES.

The following Direction was addressed to the 447 Registrars in the Country:-

"If at any time the number of deaths registered during the Quarter has been above the average, state, in a note at the foot of the Return, whether any epidemic disease, such as Measles, Typhus, &c., has been prevailing in the District; or if there be any other known circumstance which will account for the increase."

In compliance with the above direction, 202 Registrars have appended explanatory notes to their Returns; the remaining 245 Registrars apparently observed a decrease, or no obvious increase, in the mortality.

An Abstract of the Notes to the Returns is subjoined.

	***************************************	of the second se
SUP. REG. DISTRICT.	REG. DISTRICT.	
		About the average; 1 case of Typhus, and 1 of Scarlatina.
	$Yalding \dots \dots$	Small Pox has made its appearance in my district during this quarter,
		An increase on the corresponding quarter of 1843, but only I more on the total of that year. 15 were registered Natural Decay from Age, several being above 80 years: their united ages gave an average of 75 years each. 12 others died from various causes, whose united ages gave an average of 70 years each.
		The present return (70) is larger than any previous one since the passing of the Registration Act. It is 30 above the corresponding quarter of 1843; 45 more than the last quarter; and 25 above the average of three previous quarters ending in December, which may be accounted for in some measure by having a considerable influx of visitors during the past quarter, with the addition of 10 inquests. It appears that 50 were permanent residents of this district, and 20 casual.
ISLE OF WIGHT	Cowes	An increase, chiefly amongst old people above the age of 60. My district is healthy, and no Epidemic prevailing.
		In the two previous quarters I reported that Measles and Small Pox continued to increase in my district. During this quarter both Epidemics have very much abated, although 6 cases of Small Pox have proved fatal, and all of them without previous vaccination. The mortality has been greatest among the aged; 21 deaths out of 84 (the average) are between the ages of 60 and 95 years.
		I believe there has been less Epidemic and Endemic disease in my district than usual, although the number of deaths registered (108) is quite an average. The cause of death is chiefly from old age—one case being 100 years and 10 months. I have registered 5 cases of mall Pox, and all of them without previous vaccination.
		An increase of deaths, 10 more than in the corresponding quarter of 1843, chiefly among children, 12 having been registered under 2 years of age.
	St. Albans	About the average of past quarters: 26 males, 28 females. The ages of 17 of the males average 74 years.
		An increase of 17 as compared with the corresponding quarter of 1843. 8 deaths from Measles, and 5 from Typhus, all of the labouring classes.
WYCOMBE	High Wycombe	4 more than in the corresponding quarter of 1843. The increase is caused by <i>Pneumonia</i> in very young children, as no epidemic is prevailing in my district.
		quarter of 1843. 1 death from Small Pox, (natural), and 1 from Scurlating. District
		18 more deaths than in the corresponding quarter of 1843. Small Pox has been prevalent, 19 cases having proved fatal.
		The p esent quarter shews ex ctly the same number of deaths as the corresponding quarter of 1×43, and is about 5½ below the average of eight years. There are no particular diseases prevalent at this time beyond those usual in an inelement season.
CAMPRIDGE	St. Giles	Of 98 deaths registered this quarter, 6 are from Small Pox, 8 from Typhus, 2 Scarlatina,
CAMEDIAN GRAND CO.	St. Andrew the Less	About the average, and less than the corresponding quarter of 1843 by 49, which was caused by Measles and Small Pox then prevailing in my district.
COLCHESTED	St. Giles	Considerably below the every
IDSWICH	Third Ward	13 deaths by Measles this quarter.
IFSWICH	St. Margaret	Less than in any quarter since the commencement of the Registration Act, and 20 less than the corresponding quarter of 1843; while the number of births is considerably greater. The united ages of five of the persons who died amount to 400 all females.
NORWICH		Considerably above the average. Small Pox has been the prevailing epidemic, and 51 deaths have occurred; of these 2 only had been vaccinated.
******	. Mancroft	Considerably above the average, owing to the prevailing epidemics of Measles, Scarlatina, and Small Pox.
CDE ATT WATER	. Coslany	About 30 above the usual average. 17 deaths from Small Pox. Vaccination successfully performed in one case only. Meusles also prevalent.
GREAT YARMOUTH		Above the corresponding quarter of 1843, but below the number registered in the quarter e ding March, 1844. Of Typhus there are 2 cases, of Searlatina 4, and of Meusles 18. Only one death from Small Pox, and the child had not been previously vaccinated. A case is reported where death ensued, after Measles, from gangrene of the face, the result of mercurial influence.
The state of the s	. Southern	Above the average, the increase being in consequence of the prevalence of Measles amongst children, and the deaths of 14 persons over the age of 70.

_		
SUP. REG. DISTRICT	REG. DISTRICT.	W. P. C. Charlana C. L. L. M. and a control of the
DEVIZES	Devizes	My district has been particularly healthy, and the number of deaths is below the average.  Below the average. Three coroner's inquests.
DORCHESTER	Maiden Newton	Of the 24 deaths, there has been 6 sudden,—namely, 3 from Apoplexy, 2 from Disease of the Heart, and 1 from accident; also 2 children in one family from Scarlatina.
EXETER	St. David	The present return (95) is 27 less than in the corresponding quarter of 1843, although increased by 7 cases of accidental death. There are 6 deaths from <i>Measles</i> , and 6 from <i>Small Pax</i> , without previous vaccination.
	St. Sidwell	The mortality of the quarter is chiefly remarkable for Erysipelas, (which has prevailed in the hospital and in the city), and for longevity. Of the persons registered, 26 being above 60 years, 13 above 70, 8 above 80, 2 above 90, and one 107 years, 4 months, and 2 weeks.
ST. THOMAS	St. Thomas	13 less than in the corresponding quarter of 1843. Among the present number of 27 registered were eleven persons at ages varying from 67 to 88. There were 2 cases of Fever, one of Megsles, and another of Erysipelas.
	Exmouth	An increase of deaths of 12 over the corresponding quarter of 1843. 4 coroner's inquests 4 deaths from <i>Phthisis</i> , 9 of children under one year. We have no epidemic, and attribute the increase to the number of children dying soon after birth.
Section (1) (1) (1) (1) (1) (1) (1) (1) (1) (1)	Kenton	There were three inquests. Ten of the deaths were in persons beyond the natural age of man: seven at 78-9, one at 87, and one at 90.
	East Budleigh	Fever has prevailed in the district, causing 5 deaths, and 11 deaths have arisen from Consumption and disease of the lungs. One inquest on case of suicide.
Therefore Distribution ( ) ( ) ( ) ( ) ( ) ( ) ( )	Heavitree	The deaths this quarter are less than in the last, consequently the district is healthy, and yet from the variable state of the weather much sickness was looked for. There is no epidemic.
	Alphington	Below the average; the district being very healthy: 5 of the deaths were coroner's inquests.
PLYMOUTH	Saint Andrew	The mortality of this quarter is above the average, owing principally to the prevalence of fever: there have been also a few cases of Measles and Small Pox registered.
	Charles the Martyr	Several cases of Small Pox have been registered, all of which were without previous vaccination. Fever has prevailed considerably. The number of deaths is considerably above the corresponding quarter of 1843.
REDRUTH	Gwennap	Hooping-Cough and Measles have been prevalent.
	Redruth	Rather above the average, but the district is comparatively healthy.
	Illogan	Measles and Hooping-Cough appear to be the most prevalent diseases this quarter.
	Camborne	The number is 3 short of the average of the same quarter of the last 6 years.
PENZANCE	Penzance	Rather more than the average, owing to the effects of Measles and Hooping-Cough, which were also prevalent during the preceding quarter.
	St. Buryan	Two more than the corresponding quarter of 1843: the total registered by me is 15, there is but one above 20 years. My district is healthy at present.
	St. Ives	Above the average, owing to the continuance of <i>Hooping-Congh</i> and <i>Measles</i> . The number of deaths from other diseases is below an average.
Delicination and the second supplementations and the second supplementations and the second supplementations are second supplementations.	Uny Lelant	Measles, Hooping-Cough, and Typhus, have been very prevalent this quarter, and have caused an increase of 8 above the last, and 49 on the corresponding quarter of 1843, but the district appears more healthy at present.
BATH	The Abbey	A little above the average: the increase is occasioned by a larger number of deaths from Pneumonia and other diseases of the lungs.
*************	Batheaston	A little below the average.
************	Twerton	This quarter has considerably exceeded the average, but not in consequence of any epidemic. 10 have died from Consumption; Debility, 9; Hooping-Cough, 4; Convulsions, 5; and from various diseases, 21.
BRISTOL	Castle Precincts	Rather above the average, from the prevalence of fevers, diseases of the respirator organs, and accidental deaths. The deaths from Fevers, (including two cases of Typhus), are 13; Diseases of the Respiratory Organs, 27: and accidental deaths, 12 The latter number is 50 per cent. above the average.
809808	St. Augustine	A great increase in the month of December, compared with the two first months of the quarter; the numbers being respectively, 25, 24, and 40. Small Pox and Measles very prevalent: 9 cases of the former, and 11 of the latter have proved fatal.
	St. James	On comparing the returns, I find that the number is 27 above the average of the three preceding quarters, and 8 above the corresponding quarter of 1843. This may be attributed to the great prevalence of acute Pulmonary diseases, such as Pneumonia Hooping-Cough, & c., with Asthma, amounting in the whole to 27 deaths. There has been also 19 deaths from Philisis, and 4 from Measles: to these should be added 11 from Small Pox, without previous vaccination. This last lamentable fact shews plainly the inefficiency of the Vaccination Act, and that in all probability it ever will be so, unless some penal enactment is added to it, obliging people to have their children protected by vaccination within a given time. It seems to me an abuse of liberty to suffer children to be exposed to death as under the present law.
¥0580000	St. Mary Redcliff	Above the average, being 26 more than the last, and 15 more than the corresponding quarter of 1843: the number then was 110, now 125: the increase being in December, in which month I registered 61; while in October and November only 64. Hooping-Cough very generally prevailed, and proved fatal in 22 cases.
200000000000000000000000000000000000000	St. Paul	About the average. Measles and Small Pox have been prevalent: 12 fatal cases of the former, and 4 of the latter disease have been registered. The cases of Small Pox were all without previous vaccination.
CLIFTON	Clifton	mult Pox has been prevalent during the quarter: 6 deaths were registered from that disease, five without previous vaccination. The total deaths (88) is about the average number. 9 persons died at ages varying from 80 to 98.
**4******	Westbury	Diseases of the lungs have been unusually numerous during the quarter, arising from the dense state of the atmosphere, and the prevalence of Measles following Hooping-Cough. The wind during the past quarter has been with very little variation N. by N.E., standing in that direction during the dry, and wet weather: this may, with the density of the atmosphere, account for the increase in Pneumonia.
00000000	St. George	The mortality this quarter (including two inquests, both natural deaths) is 4 per cent, below the general quarterly average, and 20 per cent, below the corresponding quarter of 1843. Small Pox has prevailed extensively, and 5 deaths have occurred, all without previous vaccination.

	REG.DISTRICT.	
SUP. REG. DISTRICT.  STROUD Str		The severe and sudden changes of the atmosphere have removed many of the aged and poor
		persons, otherwise the district has been healthy and free from any prevailing diseases. The most prevailing complaints have been Consumption, Dropsy, and Hooping-Cough.  During the inclemency of the weather in the quarter 4 have died above 40, whose ages amount to 171; 3 above 60, amounting to 185; one 79, one 82, one 92: total, 253 years.
Sto	nehouse	2 have died of Fever, 2 of Scarlatina, and 1 of Brain Fever, with 2 inquests.
HEREFORD Ken	atchurch	My district has been very healthy, having registered but 17 deaths, and the greater part from old age, five being above 70 years, one 84, one 87, one 92, and one 104, a female.
SHREWSBURY St.		The deaths are 95—about 10 above the average—yet 15 less than last quarter. Total Small Pox cases 13, 8 without previous vaccination, and 5 with previous vaccination; being 13 cases of Small Pox less than last quarter. 7 other cases of epidemic or contagion have occurred, viz. Scarlatina, 3; Measles, 1; Hooping-Cough, 1; Typ us, 2. The Small Pox cases accounting for the increase above the average. There are as many as 12 persons, 3 males and 9 females, whose united ages amount to 960 years; average of age, 80 years; varying from 75 to 90 years each.
KIDDERMINSTER Kid		The number of deaths considerably exceeds the usual average. The increase arises from the great number of deaths from Fever, Typhns, Consumption, and decay of nature.
DUDLEY Ror	wley Regis	Nearly 20 more deaths this quarter than the average of the three preceding quarters. The increase is occasioned by the loss of 11 lives from an explosion of sulphuric acid gas in a coal pit, and by the prevalence of Fever in the district.
Tip	oton	The increased mortality during this quarter has been chiefly owing so the prevalence of Measles, Consumption, Bowel Complaint, Scarlatina and Typhus; of which diseases there have been registered, Measles, 23; Consumption, 19; Bowel Complaint, 12; Scarlet Fever, 11; Typhus Fever, 11; inquests, 21.
Seá	lgley	An increase of 80 deaths this quarter, the total number being 201, that of last quarter 121. The most prevalent causes appear to be Typhus, 15; (9 more than last quarter) Scarlatina, 5; Diarrheæ, 21; Enteritis, 7; Diarrheæ, Convulsions, 1; Chicken Pox and Diarrhœa, 1; Hooping-Cough and Diarrhœa, 1; Teething and Diarrhœa, 3; Measles, 2; and Measles with Convulsions, I. Gholera cases, 3, (two of which were spasmodic; Consumption, 15; Hooping-Cough, 7; Chicken Pox, 1; Thrush, 3; and 28 inquests, from various occurrences of different descriptions.
Du	dley	In this quarter the increase as compared with the last is very considerable, owing in a great measure to Fever and Bowel Complaint having been so prevalent, but more particularly the former. 47 have died of Fever, and 35 of Bowel Complaint.
		A decrease of 2 upon last quarter. 20 have died from Inflammations, chiefly of the chest and bowels, which have been the most prevailing complaints.
1		An average number of deaths. No particular complaint to notice.
		32 more than the corresponding quarter of 1843. 19 coroner's cases: 15 cases of Typhus, many of them amongst persons of a respectable order. 21 between the ages of 60 and 80, and the remainder ordinary cases. This quarter, I think, may be considered an unhealthy one.
W	illenhall	The deaths in the quarter ending December, 1843, were 51, and in the present quarter 74 shewing an increase of 23 deaths, but I am not aware that any particular epidemic has prevailed.
WOLSTANTON Tu	unstall	This district was very healthy during the months of October and November, but Measles and Inflammation of the Chest or Lungs are prevalent now.
Bu	urslem	Above the average, owing to the prevalence of <i>Measles</i> . 27 deaths took place from this cause, between November 20th, and the close of the quarter. Several deaths have been caused by <i>Croup</i> , and there were 5 cases of Premature Birth.
BIRMINGHAM St	f. George	This quarter 29 have died of Consumption, 6 of Scarlet Fever, 11 of Typhus Fever, 2 of Measles, 6 of Hooping Cough, 17 of Inflammation of the Bowels, 15 of Inflammation of the Lungs and Chest, 10 of Small Pox, without previous vaccination, and 21 inquest cases.
St	. Paul	<ul> <li>Twelve above the average of seven years; 11 have died from Small Pox, without previous vaccination, 3 from Scarlet Fever, 3 from Premature Birth, and the rest from general causes. (Two of the cases of Scarlet Fever were from one house in a filthy court.)</li> <li>Very few deaths were registered during the several weeks of severe frost.</li> </ul>
		Small Pox has been very prevalent in this district, and proved fatal in 11 cases -9 without and 2 after previous vaccination. I have every reason to believe it is now subsiding.
		No increase above the usual average. And cases of Fever producing death are reduced in number. Small Pox, without previous vaccination, 3 cases, Measles 1.
	rdington	below the average.
		An increase of 19 on the corresponding quarter of last year, which appears to have been caused by the prevalence of Scarlet Fever. of which there have been 14 cases. 6 other cases of Fever, and 4 of Small Pox.
1		• Measles have been particularly prevalent and fatal through the quarter. Out of the 82 deaths registered, Measles have been the cause in 20 cases.
	East	An increase of 16 over the preceding quarter, owing to Scarlatina still prevailing to a great extent—40 deaths from that epidemic. In other respects the district is healthy
1	West	Increase of deaths over the previous quarter 32. Scarlatina has been prevalent—24 have died, and 4 from Small Pox without previous vaccination.
LINCOLN F	Home	
	South	Typhus is at this time very prevalent in this district. Small Pox has again appeared partially, but only one death has yet been registered, and it is still confined to one part of the district.
NOTTINGHAM S	St. Ann	• 21 deaths above the average; 15 have been from Small Pox, and out of these 12 were without previous vaccination, and two of the others were only vaccinated a few days previous to their decease. A female, aged 35, died from Hypertrophy of Spleen and Liver: weight of Spleen FOUR POUNDS AND A HALF, post mortem. The surgeon who made the examination is the informant.
	Byron	· Small Pox and Measles are very prevalent.
	Arnold	One-third below the average.  Hooping-Cough has prevailed in one parish of my district, but only fatal in one case. The number of deaths returned is about the usual average, but the parasition of families (97).
		number of deaths returned is about the usual average, but the proportion of females (27 out of 39) is remarkable.

SUP. REG. DISTRICT.	REG. DISTRICT.	Annual of the state of the stat
		Various diseases have prevailed in my district this quarter, particularly Fevers and Hoop-
	Greasley	ing-Cough; the latter has in a great measure subsided.  Rather above the average, chiefly from Natural Decay, Convulsions, and Typhus.
		Precisely the same number (148) as that of the corresponding quarter of 1843.
STOCKPORT	Stockport (First)	Below the average. Quarter ending December 31st, 1843, 175; March 31st, 1844, 173; Jane 30th, 1844, 149; September 30th, 1844, 176; December 31st, 1844, 153.
	Hyde	Rather above the average, Measles prevalent. 1 death from Small Pox in a family of travelling mendicants.
		Below the average, being 42 per cent. less than its corresponding quarter; 13 per cent. less than March; 7 per cent. less than September; and only 10½ per cent. more than June. The district is particularly healthy, not having had I case of Scarlatina or Small Pox; the only fatal epidemic being Measles, which, after 7 deaths, appears to be abating. The most remarkable feature is the mortality in males, being more than 28 per cent. above that of females. Contrasting the present with its corresponding quarter in 1843, Measles is 71 per cent. more; Diarrhea and Comulsions, 60; Pneumonia, 33; Parallysis, 66; and Old Age, 100 per cent. less. Phthisis, Asthma, Apoplexy, and Fever, the same in both quarters.
	Cheadle	Total deaths 21, four of which have been coroner's inquests, 7 were at advanced ages. Total deaths in 1844, 67; in 1843, 85; in 1842, 78. 1844 is below the average.
The state of the s	Hazelgrove	The people in this district are well employed and comfortable in circumstances; consequently they are enabled to employ medical men in cases of illness, and by their instrumentality many diseases, which increased the mortality two and three years ago, are prevented from proving fatal. Considerably below the average. Influenza very prevalent, but only 2 cases have proved fatal. The Scarlatina is also prevalent, but not one case has proved fatal hitherto. There has been but one case of Typhus.
MACCLESFIELD	East	Deaths 93:—above the average, owing to the prevalence of Measles among children, and Inflammatory and Dropsical affections among the aged.
	West	Measlss has been prevalent since the middle of October, though of a remarkably mild character; 5 cases only have terminated fatally. There have been two deaths from Typhus, and one from Scarlatina.
	Bollington	15 more than the previous quarter, but not any epidemic or contagious disease apparently existing. The year ending 1844 was much below the average of deaths in 1843.
	Prestbury	Not above the average, although Small Pox has been prevalent, of which disease 1 child died, without previous vaccination; and 1 of Pneumonia, supervening after Small Pox 1 year and 9 months subsequent to vaccination.
LIVERPOOL	Great Howard Street	A decrease of 7 in comparison with last quarter. Typhus has been very prevalent here, more especially in the higher ranks. My return includes 18 cases. Small Pox appears on the decrease.
	Dale Street	58 deaths more than the previous quarter, and 36 more than the corresponding quarter of 1843, arising from cases of <i>Dropsy</i> , <i>Natural Decay</i> , Inflammation on the Lungs, Scarlatina, and 11 cases of Typhus.
The state of the s	St. Thomas	48 above the corresponding quarter of 1843. Scarlatina has been prevalent amongst children, of which I have had 42 deaths. Typhus 5, Fever 8. Diarrhea 18, Phthisis 24, Pneumonia 25, Bronchitis 16, Hydrocephalus 14, Dropsy 12, Measles 8, Teething 6, Diseased Heart 11, Abcess on the Lungs, &c., 11, Croup 3, Small Pox 2.
	Mount Pleasant	The number of deaths is 374, being more than the preceding quarter by 107, and the corresponding quarter of 1843 by 25. The most prevalent causes of death have been Typhus and Phthisis.
4	Islington	The deaths this quarter exceed those of the corresponding quarter of 1843 by 40, which increase I attribute to Asthma, Small Pox, and Old Age. I have had I2 cases of Typhus, and 10 of Scarlatina; shewing an increase of 2 in each over the corresponding quarter of 1843.
WEST DERBY	West Derby	The prevailing epidemics, arising it is presumed from the exceedingly dense state of the atmosphere, have considerably added to the mortality of this quarter. There is an increase of 46 deaths, as compared with the previous quarter.
	Everton	An increase of deaths in this quarter, arising from increase of population, from Typhus, 21. Scarlatina 5, Nervous and Brain Fever 3.
	Great Crosby	An increase of 1 over the September quarter. In this district there has been more sickness prevailing than in any preceding quarter since the commencement of the act. Tuphus and Croup were the only epidemics. Those who died of Typhus bearing the the respective ages of 19, 29, 30, and 31 years; that of Croup 5 years.
***************************************	Toxteth Park	An increase of 132 as compared with the former quarter. The prevailing diseases have been principally among children. Typhus 6, Scarlatina 62, Measles, 18, and Chronic Diarrhea 33.
BLACKBURN	Blackburn	About 100 more than the average. Small Pox has been very prevalent in my district, 117 deaths have taken place from that complaint—81 without vaccination, and 36 second attack after vaccination. Of this number about 20 were from 10 to 27 years of age. Very few of the 36 had been vaccinated by a regular surgeon: some stated that the father, others that an old woman in the neighbourhood, had vaccinated; indeed, among the lower families vaccination has been much neglected. 8 have died from Hooping-Cough, 54 from Consumption, and 29 children in fits.
		Above the usual average, in consequence of the prevalence of Scarlatina, Influenza, and Hooping-Cough, which are very severe in this district.
		Two cases of Small-pox after vaccination.
Sales Carlotte		13 more than last quarter; but the chief cause I cannot assign, as there is no epidemic disease prevailing at present in my district.
PRESTON	Preston	25 deaths less than the corresponding quarter, and 4 less than the average, although there have been 36 deaths from Small Pox.
7	Longton	30 deaths, being 6 more than in the corresponding quarter of 1843; an increase which may be attributed to the fact of more than one third being those of aged persons; 11 having occurred at ages varying from 69 to 94.
	Walton-le-dale	Under the average.
ROCHDALE	Wardleworth	About the average: the district is generally healthy, and the deaths mostly children under 2 years of age.
	Spotland nearside	My district is very healthy; there is a decrease of 5 as compared with the corresponding quarter of last year.
BURY	Bury, North	A slight decrease upon the corresponding quarter of 1843—the deaths then being 88. An increase upon the last two quarters, which were respectively 64 and 59. The deaths in October were 21, November 15, December 35—total 71.

SUP REG. DISTRICT REG. DISTRICT.						Mester !
BURY Birtle	. District in a he					191 1119
Tottington Lower End	. About an avera	ge of the corres	sponding quarters.	De la		
BOLTON Eastern	The deaths have than the aver epidemic disc died, aged 60	age of the presease. The seas years and upw	than in the correspond vious 12 months; but son of the year may per eards, one of whom wa	haps account as 103 years o	for it. 23	persons have
Western	Scarlatina, a ter from Sma death among had the diser enquiries of most of them vaccinated. neglected to	nd Small-pox, II-pox, withou est those who hase, but without the parents as have stated the In 3 cases I for avail themselves		have regis n; nor has the cinated, the second of the system of the syst	tered 10 de here been a longh many . I have a not been va fore them no tem, but had	ans this quar- , single case of of such have made (requent ccinated; and ever had them d nevertheless
Little Bolton	has ensued, a 2 of Small-pe the average.	nd 6 from Scar	orevalent, and a slight latina; 5 of Small por ation. The mortality of	of aged peop	le has been	rather above
Tamworth	No epidemic	disease has be	age, and 12 more than een prevailing in the d	in the corre	sponding qu	uarter of 1843.
WIGAN Wigan	Below the ave	rage.	13 43	mbow of 1942		
Standish		f deaths is 13 I	ess than the same qua er 8, having registered	1 38_14 ma	les and 24	females. No.
Aspull	contagious d	isease prevaili	ng at present.			
PRESCOT St. Helens	of the corres	sponding quart	t so healthy as last quers of the last four ye	ears.		
Prescot	proved fatal a fatal influe tween the ag	The severity ence on the age ges of 60 and 95	tlent in my district, by of the weather in co. During the quart years. The number	er 12 person of deaths is	s have been	registered be-
	Under the aver	age. The dea	the two former quarter ths have been amongs of deaths registered of	t old people	and very yo	ung children.
CHORLTON Hulme	part be a tri	buted to the inc	of the five previous a creased population of is have been very prev	the district.	ers. The Hooping C	excess may in ough, Meusles,
Stretford	Above the ave prevalent in latter diseas	my district; 1	ina and Hooping Cou. 4 deaths have taken p	gh have bee	n, and still he former,	continue, very and 6 from the
MANCHESTER Ancoats	mation of I Croup, 12; Convulsions 13; General Rather below Independently district are	Angs, 36; Ind Childbirth, 6 within the first Causes, 139— the average. of the Workh	ouse, Infirmary, and increase of 25 on th	vulsions with ulsions with Coroners' 1n	mption, 37 hin the firs in 1 year quests, the	; Fever, 13; it month, 12; or afterwards, deaths in my of 1843; this
London Road	excess arises  ber) by 18 year by 36, 1844, by 33 deaths from  Measles nea	s principally fring the quarter cases, than in and exceeding the unusure ulmonic April 10. Asthme	om Measles, Small Post rare 264, being more the corresponding of the average of the 7 it al inclemency of the ffections. Pneumonia of Convulsions about ent. in the correspond	than in the p parter (Dec antecedent que season has c bears a pro t 9. The su	revious qua cember) of narters, viz aused a lar oportion of abjoined tab	arter (Septem- the preceding the preceding the for 1843 and ge increase of 15 per cent.
	No. of Death	s Registered		Proportion	per cent.	Increase in
	1844	1843	Causes of Death	1844	1843	1844
	40 25 24 16 9 8	24 7 13 6 4 4	Pneumonia Measles Convulsions Asthma Hydrocephalus Scarlet Fever	15·15 9·47 9·08 6·06 3·41 3·03	10·53 3·07 5·26 2·63 1·75 1·75	4·62 6·40 3.82 3·43 1·66 1·28
	122	58		46.20	24.99	21.21
Deansgaie	last quarter that, beside tion, there! death by C to. Cases correspond The mortali before kno 250, but th 5 years of will be per rence. Th of atmospl East and w	compared with set the greater from the set of the set o	ict during the last qua the average number of nearly 100 more. 20 oges in the temperatur ery considerable, and f temperature being a The wind was chief to the amount of rain al years, being only 3.5	cting for the peral cases. any of the p. fewer in D. fewer in C. f	It is not I most part to There has receding queeember, Is a greater that the quarter oer were of eximum to the not very nied by a groints Nort quarter is to a consider	ess remarkable he infant popu- been 9 cases of arters referred 6344, than in the ann 1 have ever is about 240 or children under he minimum, it sudden occurradual increase h and South ohe smallest that able portion of
	dry season	. The mean q	eek in the quarter. I nantity of rain for the t is interesting to obse	ese months	in Manches	iter is generally

been less in amount nor mitigated in form by this remarkably dry season. The deaths from this type of diseases have been very numerous amongst the young, whilst not fewer than 32 old people above the age of 60 have died in my district during the quarter, many of them from Bronchitis affections and Congestion of the Lungs and Brain. It has fallen under my own observation and been confirmed by the reports of my brother practitioners that the Measles and Inflammatory, Affections of the Lungs, which have proved so extensively fatal to young children, have been attended by Croup of a peculiarly low type, not allowing of the active treatment usually necessary in this disease, and which has required, even from those well skilled in the diseases of children, a more than ordinary degree of discrimination and of care and attention; and when it is considered that a great majority of the cases have been attended by persons totally ignorant of the first principles of medicine, no surprise can exist at the great fatality which has occurred. The Small Pox was brought into my district by a family from a distance passing through the town, but who had taken lodgings here for a few days. Eight of the children seized with it had never been vaccinated. Great destruction to the lives of the infant population arises from their being put out to nurse, the mothers being engaged in occupations requiring their absence from home during the whole day.

Above an average; 54 deaths from Measles and Pneumonia.

SALFORD	Regent Road Greengate	Above an average; 54 deaths from Measles and Pneumonia.  Five less than the December quarter of 1843, and 21 less than the December quarter of
ASHTON	Marin College	1842. Number of deaths in October, 85; November, 85; December, 89; total, 259. Thirty per cent. above the average of the 6 corresponding quarters, the heaviest before
1000		this being 190 in 1843. There are 35 cases of Small Pox without previous vaccination, at various ages under 7 years; 3 from Scarlatina; 1 Measles; and 1 Hooping Cough. No other epidemic or endemic to notice.
	. Audenshaw & Droylsden	There are 13 deaths less than in the corresponding quarter of 1843. The district healthy, no epidemic at present prevailing.
Continues on the Continues of the Contin	. Knott Lanes	About an average of the corresponding quarters for the last three years.
	Dukinfield	An increase of 20 deaths above the corresponding quarter of 1843. 34 children have died from Convulsions, 2 from Small Pox without previous vaccination, and 12 Inquest cases.
	. Mottram	Above the average. There is no known circumstance which will account for the increase.
	Oldham above Town	Deaths 184. Deducting 30 Coroners' Inquests, (20 of which were held upon persons killed by the falling of a fire proof mill) this return presents 13 more than the average of the preceding corresponding quarters and may be accounted for by the prevalence of Typhus and Scarlet Fever; 7 fatal cases of the former, and 9 of the latter having occurred.
The second secon	. Middleton	The number exceeds the previous quarter by 12, and the corresponding quarter of 1843 by 6. The most prevailing diseases have been Consumption and Pneumonia, 10 cases of the former and 6 of the latter.
	Royton	5 more than than the average of three preceding corresponding quarters. Small-pox, Scarlatina, and Diseases of the Chest, have been prevalent. 6 cases of Small-pox, 5 without previous vaccination. Parents are culpably negligent of the means afforded them for the vaccination of their children.
	Chadderton	An unusual number of epidemic, endemic, and contagious diseases have prevailed this quarter. Hooping-cough 6, Small-pox, Dysentery, Scarlet Fever, Croup, Typhus Fever, and Bowel complaint, 1 each.
SHEFFIELD	Sheffield Park	Abone the average, apparently caused by 14 cases of Small-pox, of which number two were said to have been previously vaccinated: except two cases of Small-pox in September last, none had been registered for two years past.
	South	Above the average, which I attribute to a larger mortality than usual among the aged, occasioned by the severe weather, as there are 15 cases above 60 years of age.
	West	Rather above the average. December, 1842, 80; December, 1843, 90; this quarter, 100.  There are 16 deaths from Small-pox this quater, all previous to vaccination.
	North	Above the average by about 40, owing to the prevalence of Small-pox and Inflammation of the Lungs. Other diseases are nearly as former averages. 21 cases of Small-pox have proved fatal, all without vaccination.
800000	Attercliffe	Twelve deaths more than in the corresponding quarter of 1843. Brain Fever, Dropsy, and Phthisis, the most prevalent.
HUDDERSFIELD	Kirkburton  Newmill	An increase of 23 over the corresponding quarter of 1843, priacipally caused by Old Age—12 deaths averaged 75\frac{3}{2} each, and 22 were of infants, under 1 year. I consider the district is very healthy.  Rather above the average, but there is no particular disease prevailing in my district.
paragramma and the significant constraints of the	Honley	A little above the number registered in the corresponding quarter of 1843, namely, 32. The present number 38. But the only complaint which appears to have been unusually prevalent is Consumption, 12 having died from that cause.
	Lockwood	Somewhat above an average, though I am not aware of any epidemic prevailing at the present time.
	Halifax	The average number of deaths in the December quarter of the preceding years is 150: this quarter the number is 170, which is not a material increase considering the increase of the population.
	Southowram	One more than the corresponding quarter last year.
		22 deaths, being 5 more than the corresponding quarter of 1843. The increase is pro- bably owing to the deaths of old people.
	Sowerby	Above the average, occasioned by Measles 10, and Phthisis 17.
	Luddenden	24 deaths, being 2 more than the last quarter, but below the average of the 4 corresponding quarters of previous years. The district is tolerably healthy.
	Northowram	Rather above an average; the increase may be attributed to an unusual number of deaths from old age.
BRADFORD	East-end	One death less than in the corresponding quarter of 1843, which shews the district to have been very healthy, as the population since that time has greatly increased.
000000	Cleckheaton	About an average, and the district remains in a healthy state. There have been four Inquests this quarter.
	Manningham	Slightly above the average, no specific cause to be assigned.
		Above the average; one case of Typhus: the increase is partly children under a week
	North Bierly	19 above the average of the seven corresponding quarters; the increase has been from Phthisis, Hydrocephalus, and Inquests. Phthisis has been fatal in 13 cases, being 7 above the average; Hydrocephalus in 8 cases, 4 above the average; and Inquests 10, being 8 above the average.

SUP. REG. DISTRICT. REG. DISTRICT.	
501. 1210. 22.	Rather below an average, though less by 2 only than last quarter.
LEEDS North	Ten less than in the corresponding quarter of last year. Small-pox, Measles, and Scarlatina, have prevailed in some parts of the district, but the number of fatal cases is not large.
	Rather above an average. Scarlatina has been very prevalent—22 fatal cases.
Holbeck	The deaths exceed those of the corresponding quarter of 1843 by 19, and the average of the same quarters of the last 4 years by 34 deaths nearly. Scarlatina has been very prevalent and fatal during the quarter; one-fourth of the deaths registered are attributable to this disease alone.
Wortley	Considerably above the average. Scarlatina has prevailed, and in many cases has been fatal. It has been most malignant in the badly ventilated parts of the district, and in the low and inferior houses of the poor-
	9I deaths; 4 from Small-pox, and 3 from Typhus. Twenty-four above 50 years of age - 12 males and 12 females.
KINGSTON UPON HULL	97 deaths, including 22 cases of Small-pox, (natural), 11 Pulmonary Consumption, and 11 Convulsions, under five weeks
Mylon	Above the average. Small-pox prevailing from the 20th November 19 cases; 14 without previous vaccination, and 5 after vaccination.
YORK Micklegate	Above the average, resulting probably from a few cases of Scarlatina.
——— Walmgate	Epidemics have prevailed in my district all the year. Small-pox carried off a considerable number of children (comparatively few of whom had been vaccinated) in the spring and summer, but has now nearly disappeared. Scarlatina appeared to follow, and was very fatal during October and November. Typhus also has been prevalent since the end of September, and has proved fatal in 11 cases this quarter; Scarlatina 20, Small-pox 2. The following are the number of deaths as compared with 1842:—  1843
	March quarter       90       93         June ditto       75       115         September ditto       79       101         December ditto       92       121
Escrick	Above the average, but not from any particular epidemic disease. Out of the 16 deaths registered, the united ages of six amounts to 497 years—namely, one of 99, one of 84, one of 83, one of 80, and one of 69 years
SUNDERLAND Monkwearmouth	Somewhat above the average; the deaths chiefly occurring amongst aged people and young children.
TYNEMOUTH Tynemouth	The average number of deaths over the quarters ending, respectively, the 31st December, from the year 1837 to 1843, both inclusive, is 79: the mortality of the present quarter is 30 being only 1 above the average. In the number of deaths are twelve aged persons, who had reached above 70 years—(the greatest longevity being 90 years) the causes of deaths in most instances Natural Decay. 31 infants under 2 years of age, and five under 6 years, are comprised in the number, the causes of death being principally Hydrocephalus, Diarrhæa, Hooping-Cough, and Pneumonia. No epidemic has prevailed.
North Shields	The district is healthy; the number of deaths, 79, (of which 37 are below the age of 10 years), is about the usual average for this quarter.
Blyth	No epidemic has visited my district, and the number of deaths is comparatively few.
NEWCASTLE UPON All Saints	Hooping Cough very prevalent in my district, but hitherto not at all fatal. Deaths this quarter much below the general average, and the lowest for any winter quarter since the commencement of the Registration, except for that of 1838, which was 93.
. St. Andrew	About an average, although 17 below the corresponding quarter of last year. The district is in a very healthy state. Sof the deaths have occurred from Paralysis; average age, 61 years; four from Old Age, averaging 79 years; one from Typhus—(the only case this year)—and 32 of children under 12 years of age.
. Byker	One-third below the average of corresponding quarters in former years. 1 attribute this to the closing of two Collieries during the past year, and the removal of the families of many of the Colliers to other districts.
CARLISLE St. Mary	Twenty-one more than the preceding quarter, but about an average of the corresponding quarter of last year.
	An average return. There is no epidemic or contagious disease at present in my district.  The diseases have been principally Inflammatory, with 5 Coroner's Inquests.
	Two below the corresponding quarter of last year; 6 less than last quarter; and 16 below the number registered in the quarter ending March 31st, 1844.
	Below the average. There is no particular complaint in my district at present.  Rather above the average, Mrasles being prevalent duving the quarter, 12 deaths from
KENDALKendal	that complaint having been registered; and of the 68 deaths, 33 are under 4 years of age My district has been very healthy during the quarter; no cases of Typhus have termi-
	nated fatally. Erysipelas has spread rapidly towards the end of the quarter, but seems to be of a mild nature.
	Considerably above the average. Scarlatina and Typhus have been rather prevalent.  4 Inquests, 3 Old Age, 4 Consumption, and the remainder from various diseases.
	About the average of the corresponding quarters of the last 7 years. One-third of the number who have died this quarter (viz. 9 persons out of the 27) average the age of 75½ years each.
	An increase of eleven on the corresponding quarter of 1842, and a decrease of one on the same quarter of 1843. There is no particular disease prevailing in my district.
	The whole year has been generally healthy, and below the average.  Below the average Out of the fifteen registered there are four Coroner's Inquest cases.
MERTHYR TYDFIL . Lower	My district is at present in a healthy state.  Above the average. There are 27 cases of Measles, 29 of Phthisis, 10 of Hooping-
NEWTOWN Llanidloes, Lower	Cough, if decay from ()ld Age; but only 4 of Small-pox, which is abating very much.  Below the average. My district is in a healthy state.
WREXHAM Wrexham	About the general average. The causes of deaths are, 14, Phthiss; 9, Convulsions; 4, Dropsy; c, Old Aye; 7, Coroner's Inquests, 5 of which are accidents from explosion of fire-damp in a coal-pit: 4, Paumonia; 1, Hydrophobia, from the bite of a rabid
HOLYWELL Holywell	dog; 2, Bronchitis; 3, Hydrocephalus; and i, Small-pox, without previous vaccination. There does not appear to be any material increase if mortality as compared with the corresponding quarters since 1841, in which year it was about one-third more than those
GENERAL REGISTER OFFICE, January, 31st, 1845.	subsequently. The epidemics of Scarlatini, Pertussis, Pneumonia, and Typhus have a good deal prevailed; but Consumption has been the most fatal.
Maria Carlo	